

RESEARCH

Open Access



Horizontal jump asymmetries are associated with reduced range of motion and vertical jump performance in female soccer players

Alberto Roso-Moliner¹, Demetrio Lozano¹, Hadi Nobari^{2,3*}, Chris Bishop⁴, Antonio Carton-Llorente¹ and Elena Mainer-Pardos¹

Abstract

Background Performance in jumping and change of direction tests are good proxies to reflect the skill level during soccer-specific actions. Greater inter-leg asymmetries have been identified as a risk factor for developing acute and overuse injuries and jeopardizing soccer performance. The aim of this study was to assess the association between asymmetry in the unilateral vertical and horizontal jump tests, ankle range of motion, linear velocity, and change of direction in a sample of highly trained adult female soccer players.

Methods Thirty-eight highly trained female soccer players underwent a testing protocol including ankle dorsiflexion, single leg jumps for height (CMJ), distance (HJ), 40 m sprint and 180° change of direction tests.

Results Within-session reliability was acceptable ($CV \leq 7.9\%$), and relative reliability showed good to excellent (ICC: 0.83 to 0.99). The one-way ANOVA reported higher inter-limb differences for change of direction deficit ($10.9 \pm 8.04\%$) and single leg CMJ ($5.70 \pm 5.22\%$). Pearson correlations highlighted significant relationships between horizontal jump asymmetries and ankle dorsiflexion ($r = -0.41$), CMJ ($r = -0.36$ to -0.49) and HJ ($r = -0.28$ to -0.56).

Conclusions Assessing inter-limb asymmetries through different methods can help scientists understand the specificity of their detrimental effects on soccer performance. Practitioners should be aware of these specificities as well as the magnitude and direction of the asymmetries when aiming to improve specific on-field skills.

Keywords Symmetry, Bilateral asymmetry, Jump, ROM, Injury risk, Football

Introduction

The growth of women's soccer is a priority in FIFA's strategy, which has set a goal of doubling the number of participants between 2014 and 2026 [1]. Consequently, this growing interest has also increased research into the female game [2]. Performance in soccer requires a high level of physical fitness and mastery of sport-specific technical and tactical skills. It is an intermittent, high-intensity sport, and players must perform repeated sprints, changes of direction (COD), vertical and horizontal jumps, accelerations and decelerations [3, 4]. Consequently, these actions are often unpredictable and thus,

*Correspondence:

Hadi Nobari

hadi.nobari1@gmail.com

¹ Health Sciences Faculty, Universidad San Jorge, Autov A23 Km 299, Villanueva de Gállego, 50830 Zaragoza, Spain

² Department of Exercise Physiology, Faculty of Educational Sciences and Psychology, University of Mohaghegh Ardabili, Ardabil, Iran

³ Faculty of Sport Sciences, University of Extremadura, 10003 Cáceres, Spain

⁴ Faculty of Science and Technology, London Sport Institute, Middlesex University, London NW4 1RL, UK



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

frequently result in an uneven distribution of loading on each limb [5].

With this in mind, there has been a significant increase in studies investigating limb-to-limb asymmetry, which refers to the difference in function or performance of one limb relative to the other [6, 7]. For example, when measuring lower body asymmetries during strength tasks, exercises such as squats [8], isokinetic dynamometry [9, 10], and isometric tasks (e. g. the mid-thigh pulldown or the isometric squat) have been used [11, 12]. Although practical, some tests are expensive and time-consuming, making them difficult to use in field settings. Therefore, other methods of assessing limb-to-limb asymmetry that are less time-consuming and costly may be needed. Jump tests provide a practical and rapid form of physical assessment, often mimicking movement patterns (i.e. triple extension of lower limb joints) observed in sports [13, 14]. Previous research has used bilateral and unilateral tests to measure asymmetries, such as the horizontal jump (HJ) [15, 16] and the countermovement jump (CMJ) [15, 17]. Therefore, with a range of protocols previously used to measure asymmetry and the importance of this measure, practitioners should consider which is most appropriate for their athletes based on analysis of the needs of the sport, the players' training history and previous familiarity with the protocols. As noted, not all tests require much time, skill, or investment of money. In summary, assessing asymmetries of specific muscles or muscle groups is an important measurement that could help clarify the origin of possible asymmetries in specific performance tasks.

On the other hand, existing scientific evidence has identified ankle dorsiflexion restriction as an important risk factor for suffering acute sports-related injuries, such as anterior cruciate ligament rupture [18], as well as for developing overuse disorders such as patellar [19], and Achilles tendinopathies, shin splints [20], and anterior patellofemoral pain syndrome [20]. Consequently, it seems likely that ankle dorsiflexion range of motion (ROM) restrictions should be one of the key factors to control in this sport, as it can contribute to knee valgus momentum during landing tasks and squatting movements [20], which, in turn, increases un-wanted force demands higher up the kinetic chain at the knee [21].

In recent years, more research has examined the association between asymmetry and some physical performance tests [15, 16, 22]. For example, Maloney et al. [23] demonstrated that jump asymmetry during the unilateral jump test was associated with lower COD ($r=0.6$) performance in healthy adult males. Similarly, Bishop et al. [24] demonstrated that unilateral CMJ asymmetry of 12.5% was associated with decreased linear sprint and jump performance in youth soccer players. In contrast,

Lockie et al. [16] found no effect on COD and linear velocity performance associated with a unilateral CMJ asymmetry of 10.4% in male collegiate players.

For its part, Dos'Santos et al. [15] found an asymmetry in the unilateral HJ of 6.3% that showed no association with COD performance in a male university sample. Furthermore, to the authors' knowledge and rather importantly, few studies exist in populations of adult female soccer players [25, 26]. Of note, the cdcstudies mentioned above found jump tests are more sensitive for identifying between-limbs asymmetries (8 to 12%) than COD tests (2 to 3%) and were negative related with sprint times [25, 26]. These contradictory results indicate the need for further research to establish the relationship between these asymmetries and measures of physical performance.

Because of the above, the main objective of the present study was to demonstrate the relationship between asymmetry in the unilateral vertical and horizontal jump tests (CMJ and HJ), ankle range of motion, linear velocity, and COD in a sample of highly trained adult female soccer players. Given the paucity of data assessing the association between asymmetry and physical performance in female populations, it was challenging to elucidate a suitable hypothesis.

Materials and methods

Study design

A cross-sectional design was used in this study. Informed consent was provided to all female soccer players following the policy of their soccer club after approval by the Clinical Research Ethics Committee of the Government of Aragon (PI21/011, CEICA, Spain) and respecting the ethical standards of the Declaration of Helsinki.

Participants

Thirty-eight female soccer players (age: 23.5 ± 3.6 years; height: 163.9 ± 5.4 cm; body mass: 60.1 ± 5.5 kg; body mass index: 22.4 ± 2.3 kg/m²) participated in this study. All soccer players belonged to the same women's soccer division (Spanish Second Division) and had performed structured weekly training in terms of volume and methodology (i.e. four sessions of 90 min duration and 1 match per week) for a minimum of 10 months and had at least three years of experience in semi-professional soccer. In addition, the female soccer players were injury-free at the time of the test and during the 3-month intervention.

Procedures

Test sessions began with a rise, activate, mobilize, and potentiate system warm-up protocol [27]. Each participant was then asked to complete three unilateral CMJ and HJ preparation jumps at progressive intensities (i.e., 60%, 80% and 100% of perceived exertion) in preparation

for both tests. During the second testing session, three practice trials of a 40 m sprint and a 505 COD test were completed at the same perceived intensities. These sub-maximal trials served as a familiarization test in the field. A passive rest period of three minutes was provided before data collection. The tests were conducted the first few days of the following week of the intervention, at the same time (17:00–19:00) and under identical ambient conditions (22 °C and 20% humidity). Participants were instructed to refrain from vigorous activity for at least 48 h before data collection, and all tests were conducted on a soccer field with soccer boots.

Weight-bearing dorsiflexion ROM

The LegMotion system (LegMotion, your Motion, Albacete, Spain) was used to assess ankle weight-bearing dorsiflexion mobility (WB-DF), and the test is described elsewhere [28]. Each player performed three trials with each ankle to make the most appropriate measurement, recovering 10 s between each attempt.

Countermovement unilateral and bilateral jump

All jumps were performed on the soccer field and in soccer boots. All participants initiated each jump with hands on hips, with a leg countermovement to a depth chosen by the user before accelerating vertically and jumping as high as possible. For unilateral jumps, the leg used had to remain fully extended during the flight phase before landing on the floor. Jump height in centimetres was recorded using the “My Jump 2” smartphone app. Two jumps were performed with the right leg and two with the left leg, resting for 60 s. The highest jump was selected for further data analysis.

Standing long jump test

A standard 30-m tape measure (30 m M13; Stanley, New Britain, USA) was used. The subjects started the test standing with their feet placed behind the line and their arms relaxed. They were instructed to jump the maximum possible horizontal distance, executing a controlled landing and maintaining balance on the performing leg or both legs (3 s) until the evaluator recorded the fall position. The length was measured from the jump line to the rearmost heel in the subject’s landing. Two unilateral jump attempts were performed with each leg, and two bilateral jump attempts. Each with 60 s of recovery between jumps, the best unilateral and bilateral jump was recorded for later analysis.

40 m sprints

Photocell timing gates (Witty, Microgate, Bolzano, Italy) were placed at 0, 10, 20, 30, and 40 m. Participants started the test upright, putting their feet 30 cm

behind the starting line [29]. Two attempts were made on the soccer field, separated by a rest period of 180 s. Times were recorded in hundredths of a second. The best time was used for subsequent analysis.

505 Change-of-Direction speed test and Change-of-Direction deficit

Photocell gates (Witty, Microgate, Bolzano, Italy) were placed at 10 and 15 m. Subjects sprinted 15 m over the soccer field and performed a 180° turn with both the right and left leg, completing two trials for each leg. The time analyzed was from the 10 m photocells, the 180° turn once past the 15 m photocells, and the sprint back through the timing gates over the 10 m mark. A 180-s rest period was provided between attempts. As described by Shepard et al. [30], the best time of the right and left leg was used for further data analysis. The COD deficit (CODD) was calculated from the difference between the times of the first part of the linear sprint (i.e. time recorded in 10 m) and the COD test (i.e. 505 test) [31, 32].

Statistical analysis

SPSS statistical software (Version 25.0; SPSS Inc, Chicago, IL, USA) was used. The Shapiro–Wilk test was used to determine normality. Within-session reliability of test measures was calculated using the coefficient of variation (CV) (absolute reliability) and a two-way random intraclass correlation coefficient (ICC) (relative reliability) with the complete agreement and 95% confidence intervals. The ICC values were interpreted using Koo et al. guideline’s [33], and a CV < 10% as an acceptable responsibility criterion. It should be mentioned that asymmetries may favour either side depending on whether limb scores are higher [34]. Therefore, the consistency of the direction of asymmetry between tests was measured using the Kappa coefficient and they were interpreted as poor (≤ 0), slight (0.01–0.20), fair (0.21–0.40), moderate (0.41–0.60), substantial (0.61–0.80), almost perfect (0.81–0.99) and perfect [35]. The following equation calculated inter-limb asymmetries as the percentage difference between the two limbs [36].

$$100/\text{Max value (right and left)} * \text{Min value (right and left)} * -1 + 100$$

A one-way repeated measures ANOVA determined the systematic bias between mean asymmetry values. Pearson’s correlations were performed between inter-limb asymmetry scores and performance tests. Statistical significance was inferred from $p < 0.05$.

Results

Table 1 contains information on within-session reliability. Flexibility, jump and COD tests revealed good to excellent reliability (ICC: 0.83 to 0.99) and acceptable

Table 1 Mean scores, mean asymmetry (%) and reliability data for each test

Test ^a	Mean ± SD	Asymmetry (%)	CV (%)	ICC (95% CI)
Unilateral WB-DF (cm)				
WB-DF _R	42.8 ± 4.71	3.65 ± 3.54	2.7	0.95 (0.90–0.97)
WB-DF _L	42.02 ± 5.11		1.3	0.99 (0.98–0.99)
Unilateral CMJ (cm)				
CMJ _R	13.6 ± 1.68	5.70 ± 5.22	1.7	0.98 (0.96–0.99)
CMJ _L	13.6 ± 1.39		2.4	0.95 (0.90–0.97)
Unilateral HJ (cm)				
HJ _R	146.4 ± 12.1	2.12 ± 2.10	2.1	0.93 (0.86–0.96)
HJ _L	147.2 ± 11.3		1.9	0.94 (0.88–0.97)
Sprint				
10 m	1.93 ± 0.19	-	4	0.83 (0.70–0.91)
20 m	3.36 ± 0.21		1.9	0.88 (0.78–0.94)
30 m	4.72 ± 0.29		2.3	0.87 (0.76–0.93)
40 m	6.13 ± 0.37		2.1	0.86 (0.74–0.92)
COD				
COD _R	2.62 ± 0.16	2.99 ± 2.22	2.6	0.87 (0.76–0.93)
COD _L	2.62 ± 0.18		2.1	0.92 (0.86–0.96)
CODD _R	0.70 ± 0.24		7.9	0.91 (0.85–0.95)
CODD _L	0.69 ± 0.25	10.9 ± 8.04	7.3	0.93 (0.88–0.96)

^a ROM Range of motion, CMJ Countermovement jump, HJ Horizontal jump, COD Change of direction, R Right, L Left, SD standard deviation, CV coefficient of variation, ICC intraclass correlation coefficient, WB-DF Weight-bearing dorsiflexion test, CODD Change of direction deficit

variability (CV ≤ 7.9%). After comparing the mean asymmetry of the groups, no significant results were observed.

Table 2 shows the Pearson’s correlations between inter-limb asymmetry scores and performance tests. No significant relationships were found between CMJ and COD asymmetry and the mobility, jumping, sprinting and COD tests. On the other hand, ankle dorsiflexion ROM asymmetry was significantly correlated with CMJ right ($r = -0.36$) and 10 m sprint time ($r = 0.37$). In addition, HJ asymmetry was correlated with WB-DF ($r = -0.41$; for both legs), CMJ ($r = -0.36$; $r = -0.49$; right and left leg, respectively) and HJ right ($r = -0.56$). Finally, CODD asymmetry was statistically associated with HJ left ($r = -0.46$).

Table 3 shows the levels of agreement for the asymmetry scores (Kappa coefficient). The results showed reasonable levels of agreement between the WB-DF test and the single leg CMJ (0.29) and slight levels between single leg CMJ and HJ (0.11), CODD and single leg HJ (0.10), single leg CMJ (0.02) and WB-DF (0.02). The rest of the test’s comparison shows poor levels (-0.87 to -0.13). Individual inter-limb discrepancies are illustrated in Fig. 1 for WB-DF, HJ, CMJ, COD and CODD due to the variable nature in both the magnitude and direction of asymmetry.

Discussion

The present study examined the relationship between asymmetry in the horizontal and vertical jump, ankle dorsiflexion, linear velocity and COD in highly female soccer players. The results showed significant correlations between ankle dorsiflexion ROM and vertical jump

Table 2 Correlations between range of motions, jump and change of direction tests and asymmetry percentages

Test ^a	WB-DF Asymmetry	CMJ Asymmetry	HJ Asymmetry	COD Asymmetry	CODD Asymmetry
WB-DF _R	0.12	-0.08	-0.41**	0.05	0.08
WB-DF _L	-0.12	-0.07	-0.41**	0.16	0.14
CMJ _R	-0.36*	-0.16	-0.36*	0.04	0.08
CMJ _L	0.18	-0.15	-0.49**	0.18	0.16
HJ _R	-0.18	-0.10	-0.56**	-0.01	0.07
HJ _L	0.14	0.07	-0.28	-0.27	-0.46**
10 m	0.37*	-0.08	0.27	-0.01	0.15
20 m	-0.29	-0.04	0.19	0.12	0.23
30 m	-0.24	-0.06	0.16	0.14	0.27
40 m	-0.21	-0.03	0.17	0.18	0.29
COD _R	-0.15	-0.14	0.20	0.08	-0.01
COD _L	-0.06	-0.10	0.22	0.06	-0.08
CODD _R	0.15	-0.06	-0.09	0.18	-0.12
CODD _L	0.20	-0.11	-0.05	0.14	-0.17

^a ROM Range of motion, CMJ Countermovement jump, HJ Horizontal jump, R Right, L Left, SLROM Single leg range of motion, COD Change of direction, WB-DF Weight-bearing dorsiflexion test, CODD Change of direction deficit

Table 3 Kappa coefficients and descriptive levels of concordance of asymmetries between the jumping speed and COD tests

Test Comparison ^a	Kappa Coefficient	Descriptor
SLWB-DF -SLCMJ	0.29	Fair
SLCMJ-SLHJ	0.11	Slight
SLHJ-COD	-0.31	Fair
SLWB-DF -COD	-0.13	Poor
SLWB-DF -SLHJ	-0.21	Poor
SLCMJ-COD	-0.13	Poor
SLHJ-CODD	0.10	Slight
SLCMJ-CODD	0.02	Slight
SLWB-DF-CODD	0.02	Slight
COD-CODD	-0.87	Poor

^a SLROM Single leg range of motion, SLCMJ Single leg countermovement jump, SLHJ Single leg horizontal jump, SLWB-DF Single leg weight-bearing dorsiflexion test, COD Change of direction, CODD Change of direction deficit

performance with HJ asymmetry. The results suggest that more enormous HJ imbalances were associated with lower range of motion in ankle dorsiflexion and reduced vertical jump height. In addition, the asymmetries between ankle dorsiflexion (WB-DF), vertical jump (CMJ), horizontal jump (HJ), change of direction (COD, and CODD) rarely favored the same side, signifying the task-specific nature of asymmetry.

Table 1 shows that the metrics of inter-limb asymmetry differ depending on the test used. Therefore, female players do not all react similarly to the same asymmetry test. Literature has shown that asymmetries of > 10% may decrease jump height [37] and increase COD speed times [38], indicating that the reduction of these differences may be favourable. The first aspect to note is that higher percentages of asymmetries were identified for the CMJ than for the HJ (5.70% vs. 2.12%). Previous studies in young female soccer players [26] or recreational team sports players [16], reinforce our results, finding 12.54% and 10.4% asymmetries in the vertical jump versus 6.79% and 3.3% in the HJ, respectively. Like the authors cited above, it is thought that the vertical jump could be more sensitive than the HJ when identifying asymmetries. One explanation for this is that the movement patterns related to HJ are practised from an earlier age than those of vertical jumping [26]. In our study, the sample is composed of highly trained female soccer players, and this level of training could explain why the percentages of asymmetry in both jump tests are lower than in the other two studies cited. This is confirmed when we observe a ratio of asymmetries in the studies with adult female elite soccer players of 8.65% [39] or 7.9% in elite male players [40]. This could also explain the mean values found for the percentage of asymmetry of the ROM, which are lower (3.65%)

than those obtained in studies such as that of Madruga-Parera et al. (5.88%) [41]. The CODD follows this same trend, as our data show somewhat lower asymmetries (10.9% ± 8.04) than those found in pre-season female soccer players (24%) [42] and young players [43]. Finally, the data obtained for COD asymmetry (2.99%) are very similar to those found in other studies of adult female soccer players (2.39%) [39] or young players [43]. Further research is needed to confirm these theories regardless of the results obtained.

In Table 1, we can also observe good to excellent reliability (ICC: 0.83 to 0.99) and acceptable variability (CV ≤ 7.9%), indicating that the data can be interpreted with confidence for further analysis [44]. Again, the level of the players in our sample and their physical preparation could explain this acceptable level of variability, and the reliability obtained. Haugen et al. [45] reported a CV of 3.26% for bilateral CMJ using a force platform and a CV from 1.82% to 3.30% in 40 m linear sprints measured each 10 m. In other studies such as that of Bishop et al. [17], the CV for unilateral CMJ is 2.82% and 3.51% (right and left leg, respectively), and for HJ from 3.94% to 4.18% (right and left leg, respectively). In our case, for these tests, we obtained a CV ranging from 1.7% to 2.4% for the unilateral CMJ (right and left leg, respectively), from 2.1% to 1.9% for the unilateral SH (right and left leg, respectively), and from 1.9% to 4% for the 40 m linear sprint (measured every 10 m). In the Madruga-Parera et al. [41] study, a CV of 1.70% and 1.94% was obtained for the Ankle Dorsiflexion test and 2.88% and 2.05% for the COD test (right and left leg, respectively), results very similar to those obtained in the present study. Finally, regarding the CODD, we received a CV of 7.9% for the right leg and 7.3% for the left leg. Previous studies show results that do not differ much from those found in the present investigation (female cricketers: 8.6% for both legs; female netball players: 4.9% left leg and 6% right leg; female soccer players: 12.7% left leg and 6.6% right leg) [42]. We believe that our sample’s training and fitness level again explains the similar CV for both legs. Exell et al. [46] suggest that asymmetries can only be natural if the differences between the limbs are more significant than the variability during the test protocols. In the present study, all CV values fulfilled this criterion. Therefore, all observed differences can be considered accurate. In addition, previous studies recommend performing several measurements, calculating test variability and searching for lower CV values (acceptable < 10% or CV < 5% in unilateral jump tests), in the process of continuous monitoring of professional players [39, 47], which the authors of the present article also recommend.

Table 2 shows the Pearson correlations between the limb asymmetry scores and the measurements




Fig. 1 Individual asymmetry data for all performance tests (above 0 indicates right leg dominance and below 0 indicates left leg dominance) SLCMJ: Single leg countermovement jump; SLHJ: Single leg horizontal jump; 180° COD: 5 + 5 sprint test with a 180°; SLWB-DF: Single leg weight-bearing dorsiflexion test; CODD: Change of direction deficit

performed (unilateral WB-DF, unilateral CMJ, unilateral HJ, Linear speed 40 m measured every 10 m, COD and CODD). One of the main findings in the present study is the significant relationship between SH asymmetry and decreased performance in the CMJ right ($r=-0.36$), CMJ left ($r=-0.49$), HJ right ($r=-0.56$), WB-DF right ($r=-0.41$) and WB-DF left ($r=-0.41$). Recent publications

report that unilateral jumping performance factors (including asymmetries) are direction-specific and affect the performance of the weaker leg [26]. This assumption would partially explain the results obtained in the present work for HJ, but it does not hold for the vertical jump since the asymmetries in CMJ were not correlated with decreased unilateral jumping and cutting performance.

To our knowledge, few research have associated asymmetries between horizontal and vertical jumps. Significant correlations have been found between the percentage of asymmetry in the vertical and the HJ ($r = -0.489$; $p < 0.05$) [48]. Regarding the significant relationship between HJ asymmetry and decreased ankle dorsiflexion mobility ($r = -0.41$, both for right and left leg), we have also not found any previous research that has associated such asymmetries. However, Barrera-Domínguez et al. [49], indicate that a more significant asymmetry in dynamic balance as measured by the Y-Balance Test is associated with a lower ankle ROM and a decrease in performance in both horizontal and vertical jumping. We think all these results may be because this type of jump (e.g., HJ) is not trained as much within the specific physical preparation of female soccer players or they are not so decisive in competitive actions.

A significant relationship was also found between ankle ROM asymmetry and performance in the CMJ right ($r = -0.36$) or linear velocity measured in the first 10 m ($r = 0.37$). The study by Godinho et al. [50], shows a positive correlation between jumping ability (e.g., CMJ) and higher ankle ROM. This study also establishes a relationship between ankle dorsiflexion and dynamic balance in athletes. This aspect supports the systematic review carried out by Mesfar et al. [51], which indicates that there may be a specific association between the asymmetries of this dynamic balance and jumping performance and, on the other hand, the range of movement of the ankle and the COD tests in that involving a 180° turn. Other studies have also confirmed this, which have found significant relationships between ankle ROM and improved dynamic balance or lower body motor skills [52, 53]. About linear velocity, some studies establish a positive relationship between adequate ankle ROM and improved linear speed or centre of mass acceleration in adult sprinters [54, 55]. This provides evidence that adequate ankle dorsiflexion may be related to improved performance in the unilateral CMJ, the COD and the first few metres of the linear sprint. However, we have found little literature associated with the study of ankle ROM and performance, especially in women's soccer, and more research is needed to confirm these findings.

Finally, CODD asymmetry was significantly related to decreased HJL performance ($r = -0.46$). The literature that has analysed CODD asymmetry associated with different types of jumps is very scarce, and, in most cases, bilateral hops have been used. For example, Bishop et al. [56] analysed the % asymmetry in the CODD and compared it with different bilateral jumps (squat jump, CMJ, and drop squat). No significant relationships were obtained, but it was observed that the greater the asymmetry in the CODD, the lower the performance in the different jumps.

The most similar results to those observed in the present study have been found in a recent study [57], which has negatively associated the CODD with the jump height of the right leg ($r = -0.59$) and the concentric impulse of the same leg ($r = -0.60$). Our results would reinforce the idea of those obtained in this research and could support the concept that greater competence in the unilateral jump means better performance in the CODs. In addition, greater concentric strength could also improve this performance, as the ability to apply this type of force is decisive during the turning phase of the 505 tests.

No significant associations were found between CMJ, COD asymmetries, and the fitness mentioned above tests. These data are consistent with those presented by Lockie et al. [16] and Dos'Santos et al. [58], in which no associations were observed between CMJ asymmetry and jumping tests or time reduction in the COD (i.e., 505) in male collegiate athletes. Nor did Azahara et al. [5], find significant associations between CMJ asymmetry and total time during the agility (V-Cut) test in young male and female athletes playing team sports. The same study found significant relationships between CMJ asymmetry and 30 m linear velocity, which disagrees with our results. However, the magnitude of this relationship was small ($r = 0.26$; $p < 0.05$). Regarding adult female soccer players, Bishop et al. [39] found no correlation between vertical jump asymmetry (CMJ), 30 m linear velocity (measured at 0 m, 10 m and 30 m) and COD (505) performance. This is similar to the results shown in the study by Loturco et al. [59], in which they found no association between vertical jump asymmetries and linear velocity (30 m sprint measured every 5, 10, 20, and 30 m) and COD (zig-zag test) in adult elite female soccer players. The lack of association of vertical jump and COD asymmetries with soccer-specific performance tests in the present study may be related to the players' high level of general and specific training on this type of action [60]. In addition, a large part of the competitive actions in soccer is related to these vertical jumps or COD, which could minimise the effects of asymmetries on performance [6, 7].

Kappa coefficients were used to assess the direction of the asymmetry. Through this test, it can be observed whether the asymmetry favours one limb or the other. The results in Table 3 show fair levels of agreement between the WB-DF test and the same leg CMJ (0.29) and slight levels of agreement between the unilateral CMJ and the same-leg HJ (0.11). In addition, the one-leg CODD also showed a slight level of agreement with the same leg HJ (0.10), CMJ (0.02) and WB-DF (0.02). The rest of the test comparison shows poor levels (-0.87 to -0.13), indicating that the relationship favours the opposite leg. The individual differences between limbs are illustrated in Fig. 1, representing the magnitude and direction of the

asymmetries. Bishop [61] also suggested that the guide of the asymmetry and its importance could help to understand which limb performed better, as an absolute positive value of the asymmetry would not allow identification of the asymmetry. All of the above suggests that differences between limbs do not favour the same leg in different tests. Based on these results and previous research, the detection of asymmetries in adult female soccer players should be performed individually and using a single test for their evaluation is not recommended [6, 62].

A possible limitation of the present study is that the sample was not divided by playing position due to the small number of players. Other studies, with larger samples, divide the players by playing positions to analyse their differences [6, 62, 63]. Secondly, the Y-balance test asymmetries have been shown to be negatively correlated with CMJ height [64]. Therefore, this dynamic balance test could be included in future studies. Given that the push-off process for a COD and HJ are similar, this type of vertical jump could be added in future research because it may be more sensitive to COD performance. In addition, data cannot be extrapolated to other populations such as men's soccer, amateur level or other team sports. Despite these considerations, the present work provides relevant data on lower limb asymmetries in certain performance variables in female soccer players.

Conclusions

The present study analysed lower limb asymmetries regarding mobility, jumping and cutting in highly trained female soccer players. In summary, the asymmetries found were greater in the vertical jump and 180° COD tests. However, the asymmetries in the HJ showed the greatest agreement with the reduction in mobility, jump and COD performance. Knowing the relationships between asymmetries and performance can help coaches customize training plans for specific on-field improvements. However, the direction of asymmetry appears highly variable among female soccer players, therefore, practitioners should consider individualized analysis and including different asymmetry tests when planning an intervention program.

Abbreviations

COD	Change of direction
HJ	Horizontal jump
CMJ	Countermovement jump
CV	Coefficient of variation
ICC	Intraclass coefficient correlation
WB-DF	Ankle weight bearing dorsiflexion
ROM	Range of motion
SLROM	Single leg range of motion
SLCMJ	Single leg countermovement jump
SLHJ	Single leg horizontal jump
SLWB-DF	Single leg weight-bearing dorsiflexion test

Acknowledgements

The authors acknowledge all the participants involved in the study.

Authors' contributions

HN, CB, ACL, DL, ARM and EMP have given substantial contributions to the conception or the design of the manuscript, authors HN, ARM, and EMP to acquisition, analysis and interpretation of the data. Authors ACL, DL, and ACL have participated to drafting the manuscript, and authors HN, EMP, and CB revised it critically. All authors read and approved the final version of the manuscript.

Funding

No sources of funding were used to assist in the preparation of this article.

Availability data and materials

The data presented in this study are available on reasonable request from the corresponding author. The data are not publicly available due to privacy reasons.

Declarations

Ethics approval and consent to participate

After receiving information on the objectives and procedures of the study, participants and presents signed an informed consent form, this study complied with the ethical standards of the World Medical Association's Declaration of Helsinki (2013). The study was approved by the Clinical Research Ethics Committee of the Government of Aragon (PI21/011, CEICA, Spain). Written informed consent was obtained from the participants. Moreover, all participants were fully informed of the protocol's potential risks and benefits.

Consent for publication

N/A.

Competing interests

The authors declare no competing interests.

Received: 10 January 2023 Accepted: 3 July 2023

Published online: 08 July 2023

References

1. FIFA. Women's football member associations survey report 2019. <https://digitalhub.fifa.com/m/231330ded0bf3120/original/nq3ensohyxpuxovcovj0-pdf.pdf>.
2. Webb T, Gorczynski P, Oftadeh-Moghadam S, Grubb L. Experience and Construction of Mental Health Among English Female Football Match Officials. *Sport Psychol.* 2021;35(1):1–10.
3. Datson N, Drust B, Weston M, Jarman IH, Lisboa PJ, Gregson W. Match Physical Performance of Elite Female Soccer Players During International Competition. *J Strength Cond Res.* 2017;31(9):2379–87.
4. Datson N, Hulton A, Andersson H, Lewis T, Weston M, Drust B, Gregson W. Applied physiology of female soccer: an update. *Sports Med.* 2014;44(9):1225–40.
5. Fort-Vanmeerhaeghe A, Bishop C, Busca B, Aguilera-Castells J, Vicens-Bordas J, Gonzalo-Skok O. Inter-limb asymmetries are associated with decrements in physical performance in youth elite team sports athletes. *PLoS ONE.* 2020;15(3): e0229440.
6. Bishop C, Pereira LA, Reis VP, Read P, Turner AN, Loturco I. Comparing the magnitude and direction of asymmetry during the squat, countermovement and drop jump tests in elite youth female soccer players. *J Sports Sci.* 2020;38(11–12):1296–303.
7. Bishop C, Turner A, Maloney S, Lake J, Loturco I, Bromley T, Read P. Drop Jump Asymmetry is Associated with Reduced Sprint and Change-of-Direction Speed Performance in Adult Female Soccer Players. *Sports (Basel, Switzerland).* 2019;7(1):29.
8. Sato K, Heise GD. Influence of weight distribution asymmetry on the biomechanics of a barbell back squat. *J Strength Cond Res.* 2012;26(2):342–9.

9. Costa-Silva JRL, Detanico D, Pupo JD, De la Rocha-Freitas C. Bilateral asymmetry of knee and ankle isokinetic torque in soccer players u 20 category. *Revista Brasileira de Cineantropometria & Desempenho Humano*. 2015;17:195–204.
10. Coratella G, Beato M, Schemi F. Correlation between quadriceps and hamstrings inter-limb strength asymmetry with change of direction and sprint in U21 elite soccer-players. *Hum Mov Sci*. 2018;59:81–7.
11. Brady C, Harrison A, Flanagan E, Haff G, Comyns T. A Comparison of the Isometric Mid-Thigh Pull and Isometric Squat: Intraday Reliability, Usefulness and the Magnitude of Difference Between Tests. *Int J Sports Physiol Perform*. 2017;13:1–25.
12. Dos'Santos T, Jones PA, Comfort P, Thomas C. Effect of Different Onset Thresholds on Isometric Midhigh Pull Force-Time Variables. *J Strength Cond Res*. 2017;31(12):3463–73.
13. Bishop C, Turner A, Jarvis P, Chavda S, Read PJ. Considerations for Selecting Field-Based Strength and Power Fitness Tests to Measure Asymmetries. *J Strength Cond Res*. 2017;31(9):2635–44.
14. Loturco I, Pereira L, Kobal R, Abad C, Fernandes V, Ramirez-Campillo R, Suohomel T. Portable Force Plates: A Viable and Practical Alternative to Rapidly and Accurately Monitor Elite Sprint Performance. *Sports*. 2018;6(3):61.
15. Dos'Santos T, Thomas C, Jones PA, Comfort P. Asymmetries in Isometric Force-Time Characteristics Are Not Detrimental to Change of Direction Speed. *J Strength Cond Res*. 2018;32(2):520–7.
16. Lockie RG, Callaghan SJ, Berry SP, Cooke ER, Jordan CA, Luczo TM, Jeffriess MD. Relationship between unilateral jumping ability and asymmetry on multidirectional speed in team-sport athletes. *J Strength Cond Res*. 2014;28(12):3557–66.
17. Bishop C, Read P, McCubbine J, Turner A. Vertical and Horizontal Asymmetries are Related to Slower Sprinting and Jump Performance in Elite Youth Female Soccer Players. *J Strength Cond Res*. 2018;35(1):56–63.
18. Wahlstedt C, Rasmussen-Barr E. Anterior cruciate ligament injury and ankle dorsiflexion. *Knee Surg Sports Traumatol Arthrosc*. 2015;23(11):3202–7.
19. Hassan KA, Youssef RSE-E, Mahmoud NF, Eltagy H, El-Desouky MA. The relationship between ankle dorsiflexion range of motion, frontal plane projection angle, and patellofemoral pain syndrome. *Foot Ankle Surg*. 2022;28(8):1427–432.
20. Rabin A, Kozol Z, Finestone AS. Limited ankle dorsiflexion increases the risk for mid-portion Achilles tendinopathy in infantry recruits: a prospective cohort study. *J Foot Ankle Res*. 2014;7(1):48.
21. Halperin I, Aboodarda SJ, Button DC, Andersen LL, Behm DG. Roller massager improves range of motion of plantar flexor muscles without subsequent decreases in force parameters. *Int J Sports Phys Ther*. 2014;9(1):92–102.
22. Pardos-Mainer E, Bishop C, Gonzalo-Skok O, Nobari H, Pérez-Gómez J, Lozano D. Associations between Inter-Limb Asymmetries in Jump and Change of Direction Speed Tests and Physical Performance in Adolescent Female Soccer Players. *Int J Environ Res Public Health*. 2021;18(7):3474.
23. Maloney SJ. The Relationship Between Asymmetry and Athletic Performance: A Critical Review. *J Strength Cond Res*. 2019;33(9):2579–93.
24. Bishop C, Turner A, Read P. Effects of inter-limb asymmetries on physical and sports performance: a systematic review. *J Sports Sci*. 2018;36(10):1135–44.
25. Bishop C, Turner A, Maloney S, Lake J, Loturco I, Bromley T, Read P. Drop Jump Asymmetry is Associated with Reduced Sprint and Change-of-Direction Speed Performance in Adult Female Soccer Players. *Sports*. 2019;7(1):29.
26. Bishop C, Read P, McCubbine J, Turner A. Vertical and Horizontal Asymmetries Are Related to Slower Sprinting and Jump Performance in Elite Youth Female Soccer Players. *J Strength Cond Res*. 2021;35(1):56–63.
27. Jeffreys I. Warm up revisited—the “RAMP” method of optimising performance preparation. *UK Strength and Conditioning Association*. 2007;6:15–9.
28. Pardos-Mainer E, Casajús JA, Gonzalo-Skok O. Adolescent female soccer players' soccer-specific warm-up effects on performance and inter-limb asymmetries. *Biol Sport*. 2019;36(3):199–207.
29. Mainer-Pardos E, Gonzalo-Skok O, Nobari H, Lozano D, Pérez-Gómez J. Age-related differences in linear sprint in adolescent female soccer players. *BMC Sports Sci Med Rehabil*. 2021;13(1):97.
30. Shepard J, Dawes J, Jeffreys I, Spiteri T, Nimphius S. Broadening the view of agility: a scientific review of the literature. *J Aust Strength Cond*. 2014;22:6–30.
31. Pereira LA, Nimphius S, Kobal R, Kitamura K, Turisco LAL, Orsi RC, Cal Abad CC, Loturco I. Relationship Between Change of Direction, Speed, and Power in Male and Female National Olympic Team Handball Athletes. *J Strength Cond Res*. 2018;32(10):2987–94.
32. Nimphius S, Callaghan SJ, Spiteri T, Lockie RG. Change of Direction Deficit: A More Isolated Measure of Change of Direction Performance Than Total 505 Time. *J Strength Cond Res*. 2016;30(11):3024–32.
33. Koo TK, Li MY. A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. *J Chiropr Med*. 2016;15(2):155–63.
34. Bishop C, Lake J, Loturco I, Papadopoulos K, Turner A, Read P. Interlimb asymmetries: the need for an individual approach to data analysis. *J Strength Cond Res*. 2018;35(3):695–701.
35. Viera AJ, Garrett JM. Understanding interobserver agreement: the kappa statistic. *Fam Med*. 2005;37(5):360–3.
36. Bishop C, Read P, Lake J, Chavda S, Turner A. Inter-Limb Asymmetries: Understanding how to Calculate Differences From Bilateral and Unilateral Tests. *Strength Cond J*. 2018;40(4):1–6.
37. Bell DR, Sanfilippo JL, Binkley N, Heiderscheid BC. Lean mass asymmetry influences force and power asymmetry during jumping in collegiate athletes. *J Strength Cond Res*. 2014;28(4):884–91.
38. Hoffman JR, Ratamess NA, Klatt M, Faigenbaum AD, Kang J. Do Bilateral Power Deficits Influence Direction-Specific Movement Patterns? *Res Sports Med*. 2007;15(2):125–32.
39. Bishop C, Turner A, Maloney S, Lake J, Loturco I, Bromley T, Read P. Drop Jump Asymmetry is Associated with Reduced Sprint and Change-of-Direction Speed Performance in Adult Female Soccer Players. *Sports*. 2019;7(1):29.
40. Loturco I, Pereira LA, Kobal R, Abad CCC, Komatsu W, Cunha R, Arliani G, Ejnisman B, Pochini AC, Nakamura FY, et al. Functional Screening Tests: Interrelationships and Ability to Predict Vertical Jump Performance. *Int J Sports Med*. 2018;39(3):189–97.
41. Madruga M, Dos'Santos T, Bishop C, Turner A, Blanco D, Beltrán-Garrido JV, Perez V, Romero-Rodríguez D. Assessing Inter-Limb Asymmetries in Soccer Players: Magnitude, Direction and Association with Performance. *J Hum Kinet*. 2021;79:41–53.
42. Dos'Santos T, Thomas C, Comfort P, Jones PA. Comparison of Change of Direction Speed Performance and Asymmetries between Team-Sport Athletes: Application of Change of Direction Deficit. *Sports*. 2018;6(4):174.
43. Clemente FM, González-Fernández FT, García-Delgado G, et al. Leg dominance and performance in change of directions tests in young soccer players. *Sci Rep*. 2022;12:12900.
44. Turner A, Brazier J, Bishop C, Chavda S, Cree J, Read P. Data Analysis for Strength and Conditioning Coaches: Using Excel to Analyze Reliability, Differences, and Relationships. *Strength & Conditioning Journal*. 2015;37(1):76–83.
45. Haugen TA, Tønnessen E, Seiler S. Speed and countermovement-jump characteristics of elite female soccer players, 1995–2010. *Int J Sports Physiol Perform*. 2012;7(4):340–9.
46. Exell TA, Irwin G, Gittos MJ, Kerwin DG. Implications of intra-limb variability on asymmetry analyses. *J Sports Sci*. 2012;30(4):403–9.
47. Cormack SJ, Newton RU, McGuigan MR, Doyle TL. Reliability of measures obtained during single and repeated countermovement jumps. *Int J Sports Physiol Perform*. 2008;3(2):131–44.
48. Torreblanca Martínez V, Torreblanca Martínez S, Salazar-Martínez E. Effects of inter-limb vertical jump asymmetries on physical performance in elite soccer players under 19 years old. *J Phys Educ Sport*. 2020;20:2067–613.
49. Barrera-Domínguez FJ, Carmona-Gómez A, Tornero-Quiñones I, Sáez-Padilla J, Sierra-Robles Á, Molina-López J. Influence of Dynamic Balance on Jumping-Based Asymmetries in Team Sport: A between-Sports Comparison in Basketball and Handball Athletes. *International journal of environmental research and public health*. 2021;18(4):1866.
50. Godinho I, Pinheiro B, Junior L, Chaves G, Cavalcante J, Monteiro G, Uchôa P. Effect of Reduced Ankle Mobility on Jumping Performance in Young Athletes. *Motricidade*. 2019;15(2–3):46–51.
51. Mesfar A, Hammami R, Selmi W, Gaied-Chortane S, Duncan M, Bowman TG, Nobari H, van den Tillaar R. Effects of 8-Week In-Season Contrast Strength Training Program on Measures of Athletic Performance and

- Lower-Limb Asymmetry in Male Youth Volleyball Players. *Int J Environ Res Public Health*. 2022;19(11):6547.
52. Nakagawa TH, Petersen RS. Relationship of hip and ankle range of motion, trunk muscle endurance with knee valgus and dynamic balance in males. *Phys Ther Sport*. 2018;34:174–9.
 53. Hoch MC, McKeon PO. Normative range of weight-bearing lunge test performance asymmetry in healthy adults. *Man Ther*. 2011;16(5):516–9.
 54. Inal S, Erbuğ B, Kotzamanidis C. Sprinting, isokinetic strength, and range of motion of ankle joints in Turkish male and female national sprinters may have a relationship. *Turkish J Med Sci*. 2012;42(6):1098–104.
 55. Debaere S, Delecluse C, Aerenhouts D, Hagman F, Jonkers I. Control of propulsion and body lift during the first two stances of sprint running: a simulation study. *J Sports Sci*. 2015;33(19):2016–24.
 56. Bishop C, Clarke R, Freitas T, Arruda A, Guerriero A, Ramos M, Pereira L, Loturco I. Change-of-Direction Deficit vs. Deceleration Deficit: A Comparison of Limb Dominance and Inter-limb Asymmetry between Forwards and Backs in Elite Male Rugby Union Players. *J Sports Sci*. 2020;39(10):1088–1095.
 57. Bishop C, Berney J, Lake J, Loturco I, Blagrove R, Turner A, Read P. Bilateral Deficit During Jumping Tasks: Relationship With Speed and Change of Direction Speed Performance. 2021;35(7):1833–40.
 58. Dos'Santos T, Thomas C, Jones P, Comfort P. Asymmetries in single and triple hop are not detrimental to change of direction speed. *J Trainol*. 2017;6:35–41.
 59. Loturco I, Pereira LA, Kobal R, Abad CCC, Rosseti M, Carpes FP, Bishop C. Do asymmetry scores influence speed and power performance in elite female soccer players? *Biol Sport*. 2019;36(3):209–16.
 60. Pardos-Mainer E, Casajus J, Julian C, Bishop C, Gonzalo-Skok O. Determining the reliability and usability of change of direction speed tests in adolescent female soccer players: A systematic review. *J Sports Med Phys Fitness*. 2020;60(5):720–32.
 61. Bishop C. Interlimb Asymmetries: Are Thresholds a Usable Concept? *Strength Cond J*. 2021;43(1):32–6.
 62. Bishop C, Lake J, Loturco I, Papadopoulos K, Turner A, Read P. Interlimb Asymmetries: The Need for an Individual Approach to Data Analysis. *J Strength Cond Res*. 2021;35(3):695–701.
 63. Hammami R, Nobari H, Hanen W, Gene-Morales J, Rebai H, Colado JC, Ardigò LP. Exploring of two different equated instability resistance training programs on measure of physical fitness and lower limb asymmetry in pre-pubertal weightlifters. *BMC Sports Sci Med Rehabil*. 2023;15(1):40.
 64. Gonzalo-Skok O, Serna J, Rhea MR, Marin PJ. Relationships between Functional Movement Tests and Performance Tests in Young Elite Male Basketball Players. *Int J Sports Phys Ther*. 2015;10(5):628–38.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

