

RESEARCH

Open Access



# The physical demands and physiological responses to CrossFit®: a scoping review with evidence gap map and meta-correlation

Diogo V. Martinho<sup>1,2\*</sup>, André Rebelo<sup>3,4</sup>, Élvio R. Gouveia<sup>2,5</sup>, Adam Field<sup>6</sup>, Renato Costa<sup>1</sup>, Alex S. Ribeiro<sup>1</sup>, Juliano Casonatto<sup>7</sup>, Catarina Amorim<sup>1</sup> and Hugo Sarmento<sup>1</sup>

## Abstract

**Background** CrossFit® combines different types of activities (weightlifting, gymnastics, and cardiovascular training) that challenge aerobic and anaerobic pathways. Over the last few years, the scientific interest in CrossFit® has increased considerably. However, there have been no published reviews characterizing the physical demands and physiological responses to CrossFit®. The present study synthesizes current evidence on the physical demands and physiological responses to CrossFit®.

**Methods** The search was performed in three electronic databases (PubMed, Scopus, and Web of Science). Manuscripts related to the physical and physiological performance of adult CrossFit® participants written in English, Portuguese, and Spanish were retrieved for the analysis.

**Results** In addition, a meta-correlation was conducted to examine the predictors of CrossFit® performance. A total of 68 papers were included in the review. Physical and physiological markers differed between the different workouts analyzed. In addition, 48 to 72 h are needed to recover from a CrossFit® challenge. Specific tests that involve CrossFit® movements were more related to CrossFit® performance than non-specific.

**Conclusion** Although the characterization of CrossFit® is dependent on the workout examined, the benefits of muscle hypertrophy are aligned with the recent findings of concurrent training. The characterization of CrossFit® entire sessions and appropriate recovery strategies should be considered in future studies to help coaches manipulate and adjust the training load.

**Keywords** Physiology, Strength, Endurance, Concurrent training, Workout

\*Correspondence:  
Diogo V. Martinho  
dvmartinho92@hotmail.com

<sup>7</sup> Research Group in Physiology and Physical Activity, University of Northern Paraná, Londrina, Brazil

<sup>1</sup> University of Coimbra, Research Unit for Sport and Physical Activity, Faculty of Sport Sciences and Physical Education, Coimbra, Portugal

<sup>2</sup> Laboratory of Robotics and Engineering Systems, Interactive Technologies Institute, Funchal, Portugal

<sup>3</sup> CIDEFES, Centro de Investigação em Desporto, Educação Física e Exercício e Saúde, Universidade Lusófona, Lisbon, Portugal

<sup>4</sup> COD, Center of Sports Optimization, Sporting Clube de Portugal, Lisbon, Portugal

<sup>5</sup> Department of Physical Education and Sport, University of Madeira, Funchal, Portugal

<sup>6</sup> Department of Sport and Exercise Science, Manchester Metropolitan University, Manchester, United Kingdom



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

## Background

CrossFit® includes the training and practice of weightlifting (e.g., snatch, clean, and jerk), gymnastics (e.g., handstand walk, ring muscle pull-up), and cardiovascular activities (e.g., running, rowing, cycling) [1, 2]. The practice of CrossFit® focuses on improving different components of fitness: cardiorespiratory, stamina, strength, flexibility, power, speed, coordination, agility, balance, and accuracy [3]. Different types of workouts are prescribed: Different types of workouts are prescribed: rounds for time (RFT), performing as many rounds as possible within a given time (AMRAP), and completing repetitions of exercises in a given number of minutes (every minute EMOM). For example, the “Angie” workout consists of completing 100 pull-ups, 100 push-ups, and 100 squats as quickly as possible; the “Chelsea” workout involves performing 5 pull-ups, 10 push-ups, and 15 squats every 60 s for 30 min; and the “Nate” workout involves the completion of as many rounds as possible in 20 min of muscle-ups, handstand push-ups, and kettlebell swing [4]. Given the varied nature of CrossFit®, there are many variables (number of repetitions, sets, load, rest between sets, movement, type of workout) that can influence the response to a stimulus and, consequently, have an impact on training adaptations [2, 5].

The organization of training sessions and the adjustments of training load are central to the modality since the CrossFit® Open is a competition that allows everyone to participate in three weeks of competition. Every week, an online challenge is proposed, and CrossFit® participants should perform, record, and submit their scores [6, 7]. Subsequently, the best twenty-five percent of CrossFit® Open results advanced to the subsequent phase of competition. Then, the top forty of each region (Europe, Africa, Asia, South America, North America East, North America West, and Oceania) advance to the semifinals. After the last phase, the top 40 athletes of both sexes are selected for the CrossFit® Games [6]. Coaches organize training sessions to optimize the three essential characteristics (i.e., weightlifting, gymnastics, and cardiovascular) that constitute the CrossFit® in order to prepare the participants for ‘unknown and unknowable events [1, 2, 8]. At CrossFit® Games, athletes engage in unspecified events until right before the competition begins. The founder of CrossFit® recognized that methodological guidelines are empirical, but more evidence-based, measurable, observable, and repeatable data are needed [1].

Over the last decade, scientific interest has increased in CrossFit®, most notably in physical performance and physiology [9–11], injuries [12], psychology [13], and nutrition [14]. Reviews about CrossFit® often combine those who practice CrossFit® or are involved in functional or resistance training or those who are classified

as healthy or sedentary participants, impacting the interpretation of physical, physiological, and performance data. Moreover, previous reviews focused on physical and physiological aspects [9–11] and did not examine the demands induced by each type of CrossFit® workout (i.e., AMRAP, RFT, EMOM). Given the variability of movements, training sessions, and, consequently, the metabolic demands imposed by CrossFit® workouts, a review focused on the physical and physiological outcomes of exclusively CrossFit® participants is lacking. Another challenge is which physical protocols should be applied to predict CrossFit® performance [9]. The present scoping review aims to summarize: (1) the physical and physiological demands in CrossFit®; (2) interpret the literature that explains different activity profiles within the CrossFit® context; (3) examine the association between CrossFit® performance and physical or physiological assessments; and (4) identify literature gaps and point suggestions for further research.

## Methods

The current review followed the Cochrane instructions [15], PRISMA 2020 guidelines [16], and the respective extension for scoping reviews [16]. The protocol was developed by three expert elements and registered by an expert author on the Open Science Framework at <https://doi.org/10.37766/inplasy2024.5.0063>.

## Eligibility criteria

Original studies published in peer-reviewed journals and written in English, Portuguese, and Spanish were included in the present review. There were no defined restrictions regarding the year of publication. The inclusion criteria were defined considering the Participants, Intervention, Comparator, Outcomes, and Study Design (PICOS) framework as follows: Participants – adult CrossFit® participants with previous training experience (minimal training experience or reported training practice had to be reported for inclusion); Intervention – any outcome, observation, intervention, or exposure associated with CrossFit® participation; Comparator – optional, other sporting activities, physical active; Outcomes – physical measures (for example, performance or body composition assessments); physiological outputs (such as maximal oxygen uptake, heart rate, rate of perceived exertion, hormonal levels); Study Design – no restrictions were applied to the type of studies included in the present review. Reviews and meta-analysis were not included.

## Information sources, source strategy and selection process

Three databases were consulted (Pubmed, Scopus, and Web of Science) on 23rd February 2024 using the

following search terms: (CrossFit OR “CrossFit Games” OR “workout of the day” OR “WOD\$”) AND (“training load” OR train\* OR physiolog\* OR perform\* OR energ\* OR metabol\* OR nutrition\* OR “body composition”). Originally, the initial purpose of this review was also to include CrossFit® studies that had a nutritional component; however, a decision was taken not to include these studies in the present review since there was a wealth of data available on physical and physiological parameters in CrossFit® participants. The inclusion of nutritional studies would have reduced the focus of the current review. After extracting the papers, they were combined into a reference management software (EndNoteTM 21.0, ClarivateTM). The omissions of duplicates followed a two-step process: (1) automatically removed and (2) manually checked to ensure that duplicates were removed with precision. One author (XXX) performed the entire screening process. Two authors (XXX/XX) examined the titles and abstracts to check if the studies met the eligibility criteria. The processes were then repeated for the full texts of the included papers. When discrepancies occurred, a third author (HS) was contacted to ensure concordance by consensus.

#### Data extraction and item

The two first authors (XXX/XX) organized a predefined template to collect relevant data about physical and physiological parameters. A third author (XX) confirmed the data extraction. The final summary of data comprised information on the different papers, considering the study’s purpose and design. The data were categorized into the following topics: (1) characterization of physical (e.g., body fat percentage, lean body mass, weight lifted in CrossFit® movements) and physiological outcomes (e.g., VO<sub>2max</sub>, heart rate, blood lactate); (2) assessments of the acute effects of different CrossFit® workouts on physical performance measurements; (3) data pertaining to the chronic effects of CrossFit® participation; (4) comparisons of CrossFit® participation with other types of training or no training; (5) interventional studies with participants of different competitive levels; (6) comparisons of different competitive levels in physical and physiological outcomes; (7) predictors of CrossFit® performance. Corresponding authors were individually contacted when the information was unavailable. In case the authors did not respond and the data were presented graphically, a specific software was used (GetData Graph Digitizer; <http://www.getdata-graph-digitizer.com>). This software has been shown to be accurate and precise [17] to extract mean and standard deviation from graphs.


#### Statistical analysis

Data analyses were conducted using Comprehensive Meta-Analysis software (CMA, version 2.2.064, Biostat, NJ, USA). The main outcome was performance, which was divided considering three different groups: (1) global (overall performance or ranking), (2) studies focused on a challenge from a CrossFit® competition, only reported the challenge as workout of the day, rounds for time or as many rounds as possible, or combined different challenges, or (3) studies using particular workouts (e.g., Donkey Kong, Fran, Grace, Murph, Nancy). Different protocols were used to examine the relationship with CrossFit® performance and were classified according to the methodologies reported: specific from CrossFit® (e.g., snatch, bench press, squat) and non-specific (e.g., VO<sub>2max</sub>, Wingate test). The statistical method for examining the relationship between two continuous variables over numerous studies is meta-correlation, also known as meta-analysis of *r* correlation coefficient. From each study, the coefficient of correlation and the number of participants were retrieved. The magnitude of correlation coefficients was interpreted as follows [18]: trivial (*r* < 0.10), small (0.10 ≤ *r* < 0.30), moderate (0.30 ≤ *r* < 0.50), large (0.50 ≤ *r* < 0.70), very large (0.70 ≤ *r* < 0.90), and nearly perfect (*r* ≥ 0.90). The I<sup>2</sup> informed about the proportion of variance in correlation coefficients that was due to heterogeneity instead of chance. The cut-off values for I<sup>2</sup> values were interpreted as follows [19]: low (I<sup>2</sup> < 25%), moderate (25% ≥ I<sup>2</sup> > 75%), and high (I<sup>2</sup> ≥ 75%). Subgroup differences between different types of protocols were tested, and statistical significance was determined at a two-sided level of *p* < 0.05.

## Results

#### Study identification and selection

The initial search was conducted in three databases, and 2238 records were detected. Of these, 818 were identified as duplicates, and once removed, 1520 papers were screened according to title and abstract. This process resulted in full-text screenings of 158 studies. Nine reasons were identified to exclude 117 studies: nutrition (*n* = 38), training experience (i.e., average or minimal experience) (*n* = 54), not an original paper (*n* = 8), not physical or physiological outputs (*n* = 6), papers about injuries (*n* = 5), combined different sports activities (*n* = 2), studies that examined music effects on performance (*n* = 2), detraining (*n* = 1), and psychological outputs (*n* = 1). Finally, 68 studies were included in the current review and were retrieved for the analysis, as shown in Fig. 1.



\*Since the number of papers for physical and physiological parameters was considerable, authors will consider the relevant nutritional aspects of Crossfit® participants in a separate review.

**Fig. 1** Prisma flow diagram of the study identification and selection to the present review

### Study characteristics

Table 1 summarizes the primary information extracted from each study, the origin of the corresponding author, the inclusion criteria to be considered as a CrossFit® participant, sampling characteristics, aim, main methodologies or variables analyzed, as well as findings of each study [10, 20–86]. Twelve studies focused on the comparison between CrossFit® workouts or contrasted CrossFit® with types of training [22, 24, 25, 35, 41, 45, 50, 54, 63, 68, 74, 85]. Five studies distinguished participants of CrossFit® across different competitive levels [27, 31, 34, 43, 84]. Five studies focused on recovery [20, 38, 43, 71, 83] while seven papers centered on testing the data quality of protocols that can be useful for CrossFit® [32, 44, 46–48, 55, 75]. More than 50% of papers ( $n=38$ ) focused on the physical and physiological characterization of CrossFit® participants. Twenty-three (~34%) of the studies were organized by Brazilian authors, and American authors led 19 studies (~28%). Among European countries, Spanish authors were interested in research in CrossFit® ( $n=12$  manuscripts, ~18%) – Fig. 2, panel A. Data regarding the training experience in CrossFit® varied across the studies (Fig. 2, panel B). Most of the studies

included participants with at least one year of experience ( $n=24$ , ~35%), and the 6-month cut-off value to describe training experience was used in 15 studies (~22%). Three years of training experience was uniquely considered in four studies ( $n=4$ ).

### Results of individual studies

As shown in Fig. 3 (panel A), regarding the evaluation of maximal oxygen uptake (i.e.,  $\text{VO}_{2\text{max}}$ ) during different workouts (i.e., Fran, Isabel, Cindy), the values varied within the same training workout and across metabolic challenges. Considering the Fran workout exclusively, the  $\text{VO}_{2\text{max}}$  reported in 20 participants with more than three years of practice was  $49.2 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  [20], whilst the value was substantially lower in 10 participants with one year of CrossFit® experience ( $29.1 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ) [85]. The  $\text{VO}_{2\text{max}}$  of other metabolic challenges (i.e., Isabel and Cindy) was also substantially different than the values obtained in response to the Fran Workout [21, 85]. The overall mean of maximal heart combining the data points was 184 beats per minute, with individual values of each study ranging from 177 beats per minute on an as many rounds as possible workout [87] to 189 on rounds

**Table 1** Summary of sampling, aim, methodologies and results of studies that investigated physical and physiological dimensions of CrossFit® participants

| Study                | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics  | Aim   | Main methodologies/ variables extracted  | Main results   | Main topic   |
|----------------------|---------------------------|---|---|---|--|--|--|
| Rios et al. [20]     | Portugal                  | ≥3 yrs                                  | N=14 males, N=6 females (age: 29.6±4.0 yrs, training experience: 5.9±1.1 yrs) | Describe the physical impact of Fran workout and recovery period  | VO <sub>2max</sub> , heart rate, lactate, RPE, jumping, force, velocity            | After 24 h of the CrossFit® workout, the values of jumping outputs and force are ↓ compared to the baseline.   | Physical and physiological characterization; Physical – recovery |
| Rios et al. [21]     | Portugal                  | ≥3 yrs                                  | N=14 males (age: 28.3±5.4 yrs, training experience: 5.6±1.8 yrs)              | Describe the physiological impact of Isabel workout and recovery period   | VO <sub>2max</sub> , RER, HR, lactate, RPE   | ↑ RER after 5 min to finish the workout<br>↓ HR decreased from 177 to 128 bpm 5 min after to stop the exercise   | Physiological characterization                                   |
| Carvalho et al. [22] | Brazil                    | ≥2 months                               | N=7 males, N=22 females (age: 29.7±5.2 yrs, training experience: NR)          | Compare two different types of training (running vs. CrossFit®) on anthropometric characteristics, cardiorespiratory fitness, sleep quality and lipids parameters | Pittsburgh Sleep Quality Index, cardiorespiratory fitness and lipid profile        | ↓ volume of training in running training<br>↑ VO <sub>2max</sub> running training  | Comparison between workouts                                      |
| Santos et al. [23]   | Brazil                    | >6 months                               | N=21 males (age: 26.4±4.1 yrs, training experience: 2.1±4.1 yrs)              | Compare physiological outputs on CrossFit® training and high intensity continuous training  | HR, lactate, RPE, RPD  | ↑ HR <sub>max</sub> lactate, RPE and RPD in CrossFit® training<br>↔ HR <sub>mean</sub> in both trainings   | Comparison between workouts                                      |
| Rios et al. [24]     | Portugal                  | ≥3 yrs                                  | N=16 males, N=4 females (age: 26.0±5.0 yrs, training experience: 5.8±2.2 yrs) | Characterize the Fran workout according to metabolic pathways   | VO <sub>2max</sub> , HR, RPE   | During Fran:<br>↓ aerobic (29–52%) and anaerobic alactic (23–30%)<br>↑ lactate (18–48%)  | Physiological characterization                                   |
| Pearson et al. [25]  | U.S                       | >1 yr                                   | N=13 males, N=8 females (age: 18–30 yrs, training experience: NR)             | Compare aerobic capacity, metabolic response, mitochondrial capacity, large vessel function and between CrossFit® participants and control group                  | VO <sub>2max</sub> , body fat, lactate, mitochondrial and vascular characteristics | ↑ VO <sub>2max</sub> in CrossFit® participants<br>↑ post lactate exercise in CrossFit® participants<br>↓ body fat (CrossFit® group: 18.6±3.8%, control group: 30.3±8.4)<br>CrossFit® participation ↑ mitochondrial and vascular function | Physiological characterization                                   |

**Table 1** (continued)

| Study                        | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics   | Aim   | Main methodologies/variables extracted  | Main results   | Main topic                      |
|------------------------------|---------------------------|---|--|---|---|--|---------------------------------|
| Párraga-Montilla et al. [26] | Spain                     | >6 months                               | N=72 males, N=18 female<br>(age males: 33.2±6.0 yrs, age females: 30.1±6.9 yrs, training experience: NR) | Describe neuromuscular characteristics considering the force velocity profile                   | Squat jump, countermovement jump, drop jump   | ↑ velocity than force which highlights deficits in force–velocity profile  | Physical characterization       |
| Meier et al. [27]            | Germany                   | NR                                      | N=27 males and females (age: 30.9±4.2 yrs, training experience: 1.3±1.1 yrs)                             | Examine differences on physiological parameters using athletes of contrasting experience levels | HR  | Variation in maximal heart rate within CrossFit® session<br>↔ no differences in HR <sub>mean</sub> and HR <sub>mean</sub> between contrasting groups experience<br>↔ body fat percentage between competitive levels                                    | Comparison by competitive level |
| Manrique et al. [28]         | Colombia                  | >1 yr                                   | N=30 males (age: 26.6±5.2 yrs, training experience: NR)  | Compare physical performance of CrossFit® athletes of contrasting experience athletes           | Body fat, RM squat, RM bench press  | RM bench press and squat ↑ in elite group  | Comparison by competitive level |
| Mangine et al. [29]          | US                        | Participation in CrossFit® Open         | N=80 males, N=80 females (age: NR, training experience: NR)  | Test the impact of sex and rank on pacing strategy during the 2020 CrossFit® Open               | Repetition per test, repetition per minute, failed repetitions, break times, transition times | Top male participants were 17.5% faster during tests 1, 3 and 5<br>Top female participants were 9.5% faster in tests 1 and 3<br>Top 10% athletes were faster<br>Males keep their pace during resistance exercises in comparison to female participants | Physical characterization       |

**Table 1** (continued)

| Study                               | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics   | Aim  | Main methodologies/variables extracted  | Main results   | Main topic                                  |
|-------------------------------------|---------------------------|---|--|--|---|--|---|
| Mangine et al. [30]                 | U.S                       | Participation in CrossFit® Open         | N=476,346 males, N=216,261 females (age: NR, training experience: NR)              | Creating percentiles for CrossFit® Open, examined differences between male and females     | Workout performances  | Differences between sexes were noted for most of the workouts (<93%) Workouts scored by repetitions completed—↑ repetitions in 18 workouts in 6 workouts among females                               | Physical characterization                   |
| Mangine et al. [31]                 | U.S                       | Participation in CrossFit® Open         | N=550,000 males, N=550,000 females (age: NR, training experience: NR)              | Test performance variation and sex variation in workouts and subsequent impact on ranking  | Workout performances  | Workouts assessed by time – males faster in 10 workouts whilst females were faster in 6 workouts Males lifted ↑ load in three workouts   | Physical characterization                   |
| Linhares et al. [32]                | Brazil                    | NR                                      | N=35 males (age: 31.0±5.2 yrs, training experience: 3.2±1.6 yrs)                   | Test the association between strength and power with power clean movement                  | Isometric mid-thigh pull, countermovement jump, RM power clean                  | Maximal isometric mid-thigh pull was positively related RM power clean ( $r=.51, p<0.05$ ) Lower limbs power output and rate of force development were weakly related to the power clean performance | Testing                                     |
| da Silveira Castanheira et al. [33] | Brazil                    | >2 yrs                                  | N=16 males and females (age: 31.0±6.8 yrs, training experience: 2.8±0.8 yrs)       | Examine physical and physiological performance across three months of CrossFit® training   | Sleep quality, pain score, recovery index, countermovement jump, HR variability | ↔ sleep quality, countermovement jump and RMSSD Fluctuations on pain score were found  | Physical and physiological characterization |
| Britto et al. [34]                  | Brazil                    | >1 yr                                   | N=32 males and females (age and training experience reported by competitive level) | Test the impact of FRAN WOD on psychophysiological outputs in different competitive levels | Cognition parameters, HR, blood pressure  | Changes in cognitive levels were noted in all competitive groups<br>↔ HR and blood pressure on post-test of elite and beginner participants  | Comparison by competitive level             |

**Table 1** (continued)

| Study                      | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics   | Aim  | Main methodologies/variables extracted   | Main results   | Main topic                         |
|----------------------------|---------------------------|---|--|--|--|--|------------------------------------|
| Barreto et al. [35]        | Brazil                    | >1 yr                                   | N=9 males (age: 29.6±3.5 yrs, training experience: NR)   | Test the impact of different workouts in heart rate and blood pressure   | HR variability, blood pressure   | After workouts ↓ parasympathetic indexes and ↑ sympathetic indexes<br>After WODs ↓ in systolic blood pressure  | Comparison between workouts        |
| Schlegel et al. [36]       | Czech Republic            | Qualifying for the CrossFit Games®      | N=40 males and females (age males: 27.2±3.7 yrs, age females: 27.8±5.1 yrs, training experience: NR) | Compare sex performance in CrossFit® competitions  | Workout performances   | In most of events, males attained better performance than females (difference ranged from 0.1 to 33.1%)<br>The biggest difference was in RM of snatch  | Physical characterization          |
| Menargues et al. [37]      | Spain                     | >2 yrs                                  | N=19 males, N=8 females (age males: 39 yrs, age females: 28 yrs, training experience: NR)            | Describe anthropometric profile of athletes and the impact on physical performance                                       | Body composition (anthropometry), workout performances   | A positive and a significant relationship between muscle mass and total load performed ( $r=0.876$ ) in Total CrossFit® workout  | Physical characterization          |
| Martinez-Gomez et al. [38] | Spain                     | >1 yr                                   | N=15 males (age: 29.0±8.0 yrs)   | Compare different methods of recovery (cycling, neuromuscular electrical stimulation, control) in CrossFit® participants | RPE, delayed-onset muscle soreness, HR blood lactate, muscle saturation, counter-movement jump | ↓ RPE in neuromuscular compared to control group<br>Negligible differences were found between the remaining protocols  | Physical and physiology – recovery |
| Mangine et al. [39]        | U.S                       | >1 yr                                   | 19 male participants (age: 31.0±6.8 yrs, training experience: NR)                                    | Test the association between body composition and FRAN performance   | Body composition (dual-energy X-ray absorptiometry)  | Body composition was related to performance in male and females in FRAN performance  | Physical characterization          |
| Mangine et al. [40]        | U.S                       | >1 yr                                   | 220 male CrossFit® Open athletes (age: 28.5±4.4 yrs, training experience: NR)                        | Test the influence of previous performance on 2020 CrossFit® Open results  | Workout performances   | The association is also modulated by competitive level and rank<br>Previous ranking and regional appearances were associated with overall and weekly subsequent performance in 2020 CrossFit® Open | Physical characterization          |

**Table 1** (continued)

| Study                  | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics  | Aim   | Main methodologies/variables extracted   | Main results  | Main topic                 |
|------------------------|---------------------------|---|---|---|--|---|----------------------------|
| Forte et al. [41]      | Brazil                    | >6 months                               | N=11 males, N=12 females (age: 25.9±3.6 yrs, training experience: NR) | Compare the physiological performance in two different workouts (Cindy and Open Access 18.4)    | HR, blood pressure, lactate  | Physiological differences were noted between Cindy and Open Access 18.4 workouts  | Compare different workouts |
| Dias et al. [42]       | Brazil                    | >6 months                               | N=15 males (age: 26.0±6.5 yrs, training experience: 1.2±0.3 yrs)      | Examine acute training load in CrossFit® sessions   | HR, RPE  | HR ↑ mobility and workout segments<br>HR was, on average, 65.1±5.4% HR <sub>max</sub><br>RPE and HR ↑ increased in each segment of training   | Physical characterization  |
| Sousa-Neto et al. [43] | Brazil                    | >6 months                               | N=8 males (age: 28.4±6.4 yrs, training experience: NR)                | Determine the recovery time of Karen workout  | Countermovement jump, CK, PRS  | ↑ CK 24-h after the baseline jump<br>↓ PRS decreased 24 h after exercise and ↑ in subsequent hours  | Physical recovery          |
| Conde et al. [44]      | Brazil                    | >10 months                              | N=7 females (age: 18–40 yrs, training experience: 1.9±1.2 yrs)        | Examine the sensitivity (changing detection) of physical tests and subjective scale of recovery | Perception of recovery (subjective scale), Karen performance, repetitions until failure (squat, bench press), countermovement jump, tapping test | Two microcycles were examined (aerobic and strength); in strength microcycle the squat resistance test was associated with training load ( $r=0.76$ ). In opposition, any physical test was associated with training load in aerobic microcycle | Testing                    |

**Table 1** (continued)

| Study                    | Country of contact author | Inclusion criteria (CrossFit® practice)          | Sample characteristics   | Aim  | Main methodologies/variables extracted  | Main results   | Main topic                 |
|--------------------------|---------------------------|--|--|--|---|--|----------------------------|
| Toledo et al. [45]       | U.S                       | >1 yr  | N=13 males, N=10 females (age: 26.5±4.3 yrs, training experience: 2.0±0.6 yrs) | Compare physiological outputs in two different types of training (i.e. as many rounds as possible and rounds for time) | HR, blood lactate, RPE  | Among females:<br>↔ HR <sub>max</sub> and lactate<br>↑ RPE in both type of workouts for time workout   | Compare different workouts |
| Tibana et al. [46]       | Brazil                    | >6 months  | N=11 males, N=6 females  | Assessed the relationship between physical performance and CrossFit® Open 2020   | Body composition (dual-energy X-ray absorptiometry), VO <sub>2max</sub> strength, power, muscular endurance                   | CrossFit® specific tests (strength, endurance) had ↑ association with performance<br>Negligible associations between performance, body fat and cardiorespiratory fitness were noted  | Physical characterization  |
| Schlegel et al. [47]     | Czech Republic            | Best results in the Czech CrossFit® Open ranking | N=20 males (age: 28.5 years yrs, training experience: 5.1 yrs)                 | Test the relationship between strength and endurance with CrossFit® Open   | Questionnaire about current performance   | Olympic movements (snatch, clean and jerk) were associated with classification in the CrossFit® Open   | Testing                    |
| Ponce-Garcia et al. [48] | Spain                     | >1 yr  | N=19 (age: 28.6±6.6 yrs, training experience: NR)                              | Compare laboratory and field protocols   | Anaerobic squat tests (60% of body weight, 70% of body weight), repeated jump test, assault bike test, Wingate anaerobic test | Compared to Wingate anaerobic test, a systematic bias of field test ranged from -110 to 464 watts<br>Differences between protocols were significant                                  | Testing                    |
| Peña et al. [49]         | Spain                     | >2 yrs   | N=10 males (age: 28.8±3.5 yrs, training experience: NR)                        | Test the relationship between anthropometric, physical fitness and performance   | Anthropometric measurements and fitness tests   | CrossFit® performance was associated with snatch load (corresponding to the ↑ mean power)<br>The combination of different physical tests also explained 72% of CrossFit® performance | Physical characterization  |
| Mota et al. [50]         | Brazil                    | >1 yr  | N=10 males (age: 30.0±6.6 yrs, training experience: NR)                        | Analyze the effects of two CrossFit® protocols on glycemia level   | Glycemia level  | ↑ glycemia levels in Diana and Cindy workouts compared to pre-test<br>↔ glycemia in both workouts  | Compare different workouts |

**Table 1** (continued)

| Study                        | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics  | Aim  | Main methodologies/variables extracted  | Main results  | Main topic                                  |
|------------------------------|---------------------------|---|---|--|---|---|---|
| Meier et al. [51]            | Germany                   | NR                                      | N=66 male, N=96 females (age: 32.6±8.2 yrs, training experience: 3.4±1.9 yrs)                             | Define athletic patterns for American and German CrossFit® athletes            | Questionnaire about benchmark profile   | Significant correlations were found between the power lift and Olympic lifts ↑ values of explanation between RM squat and Olympic lifts (snatch, clean and jerk)  | Physical characterization                   |
| Mangine et al. [52]          | U.S.                      | >6 months                               | N=5 males, N=6 females (age males: 34.4±3.8 yrs, age females: 35.2±6.3 yrs, training experience: 1–5 yrs) | Evaluate pacing strategies in CrossFit® athletes and the impact on performance | Workouts performance and pacing variables (repetition rate)                         | Average rate of round predicted ≥89% of performance in difference workouts<br>Other variables were determinant to explain performance in different workouts; competitive level, rest between thrusters and burpees<br>↑ blood lactate, HR, RPE in 2 km rowing and maximal repetition of thrusters | Physical characterization                   |
| Letiao et al. [53]           | Portugal                  | >3 yrs                                  | N=15 males (age: 24.0±4.2 yrs, training experience: NR)   | Determine the physical and physiological impact of performance in FRAN WORD    | RM pull-ups, RM and thrusters, row test, FRAN performance                           | FRAN performance was associated with RM of thrusters and pull-ups, RM thrusters, 2 km rowing No differences were found between groups   | Physical and physiological characterization |
| Fernando et al. [54]         | Brazil                    | >6 months                               | N=16 males (age: 29.0±8.5 yrs, training experience: 1.3±0.9 yrs)  | Compared two different types of training (CrossFit® vs. Cross-training)        | Body composition (anthropometry) and physical tests                                 | No differences between groups were found between performance, metabolic markers and hormones  | Physical characterization                   |
| Fernández-Lazaro et al. [55] | Spain                     | >1 yr                                   | N=10 males (age: 38.4±3.8 yrs, training experience: NR)   | Examine the effects of elevation training mask (12 weeks)                      | CrossFit® performance, lactate dehydrogenase, CK, myoglobin, testosterone, cortisol | No differences between groups were found on performance, testing  | Physical characterization                   |
| Bustos-Viviecas et al. [56]  | Colombia                  | > 10 months                             | N=4 males, N=4 females  | Test the association between body fat and physical performance                 | Body composition (anthropometry), cardiorespiratory fitness                         | Robust associations between body fat percentage and cardiorespiratory fitness were found in males ( $r=-0.94$ ), females ( $r=-0.95$ ) and whole sample ( $r=-0.87$ )   | Physical characterization                   |

**Table 1** (continued)

| Study                      | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics  | Aim  | Main methodologies/variables extracted  | Main results  | Main topic                                  |
|----------------------------|---------------------------|---|---|--|---|---|---|
| Pritchard et al. [57]      | New Zealand               | Participation on 2018 CrossFit® games   | N=33 males, N=39 females  | Describe tapering practices in CrossFit® athletes          | Online survey   | Most of the athletes (~99%) tapered preceding important competitions 90% of athletes ↓ training duration  | Physical characterization                   |
| Martínez-Gómez et al. [58] | Spain                     | > 1 yr                                  | N=15 males (age: 35.0±9.0 yrs, training experience: 3.3±2.2 yrs)        | Identify the main predictors of CrossFit® performance      | Physical (squat, bench press, jumping), laboratory tests (treadmill and Wingate), CrossFit® performance   | The associations between CrossFit® performance and jumping performance, Wingate test, relative strength (bench press and squat), $\dot{V}O_{2\text{max}}$ and speed were large (r value ranged from 0.58–0.75).   | Physical and physiological characterization |
| Mangine et al. [59]        | U.S.                      | > 1 yr                                  | N=8 males, N=8 females (age: 30.7±6.9 yrs, training experience: >2 yrs) | Identify the main predictors of CrossFit® Open performance | Resting energy expenditure, hormone parameters, body composition (4-compartment model), muscle morphology (ultrasound), cardiorespiratory fitness, isometric strength, CrossFit® Open performance | Body fat percentage was determinant to explain performance in three workouts ( $R^2$ ranged from 0.55 to 0.89). Other predictors also explained considerably the CrossFit® performance: training experience, muscle morphology, cardiorespiratory parameters and rate of strength development | Physical and physiological characterization |

**Table 1** (continued)

| Study                     | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics   | Aim   | Main methodologies/variables extracted  | Main results  | Main topic                      |
|---------------------------|---------------------------|---|--|---|---|---|---------------------------------|
| Mangine et al. [60]       | U.S                       | >2 yrs                                  | N=16 males and females (experienced group – age: 27.8±4.2 years, training experience: 6.4±5.6 yrs; recreational group: 33.5±8.1 yrs, training experience: 3.3±1.7 yrs) | Examine differences in anthropometric, hormonal and physiological of participants in contrasting competitive levels (advanced, recreational, control) | Resting energy expenditure, body morphology, aerobic condition, strength, blood samples   | Advanced participants ↓ body fat percentage, ↑ muscle morphology, strength, aerobic capacity and 3-min maximal test ↪ recreational and control groups have identical characteristics  | Comparison by competitive level |
| Gomez-Landero et al. [61] | Spain                     | >2 yrs                                  | N=15 males (age: 30.5±5.5 yrs, training experience: NR)  | Test the association between morphological variables and performance  | Skinfolds, RM squat, maximal, RM bench press, maximal pull-ups, sit-ups (60 s), countermovement jump, shuttle run test ( $\dot{V}O_{2\max}$ ), CrossFit® performance (Fran and Donkey Kong) | Donkey Kong was related with $\dot{V}O_{2\max}$ ( $r=-0.68$ ), suprailiac skinfold ( $r=0.71$ ) and sit-ups ( $r=-0.56$ ) Fran workout was related with squat performance ( $r=-0.53$ )   | Physical characterization       |
| Gomes et al. [62]         | Brazil                    | >3 months                               | N=23 (age: 31.0±1.0 yrs, training experience – novice: 0.5±0.1 yrs, experienced: 2.4±0.2 yrs)  | Examine the acute impact of CINDY workout regarding muscular markers, inflammatory system and stress  | White blood cell, CK, cortisol, lactate   | Post exercise noted ↑ white blood cells, lymphocyte, cortisol in experience athletes<br>↔ CK levels were found in different competitive levels  | Physiological characterization  |
| Failli et al. [63]        | Italy                     | 1 yr                                    | N=20 males (age – CrossFit® group: 26.4±3.4 yrs, RT: 26.3±3.6 yrs, training experience: 1 yr)  | Compare acute and catabolic markers in two different types of training (CrossFit® vs. resistance training)  | Cortisol, interleukin 1-beta, uric acid   | Acute effects of CrossFit® training ↑ cortisol levels whilst, in resistance training the cortisol levels ↓ Chronic effects ↓ cortisol levels in CrossFit® training and negligible changes were noted for resistance training<br>IL-1 $\beta$ ↓ after CrossFit® and resistance training (acute effects)<br>Uric acid ↑ after CrossFit® and resistance training (acute effects) | Compare different workouts      |

**Table 1** (continued)

| Study                 | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics  | Aim   | Main methodologies/variables extracted  | Main results  | Main topic                                  |
|-----------------------|---------------------------|---|---|---|---|---|---|
| Dexheimer et al. [64] | U.S                       | >1 yr                                   | N=17 males (age: 29.5±5.6 yrs, training experience: NR)   | Test the impact of muscular strength in workout performance   | RM back-squat, strict shoulder press, deadlift, workouts performance. The combination of RM was defined as total body strength                                    | Total body strength explained 62% of variance on 30 clean and jerks time performance<br>No significant correlations between workouts and total body strength were found                           | Physical characterization                   |
| Cavedon et al. [65]   | Italy                     | >1 yr                                   | N=24 males (age: 28.2±3.4 yrs, training experience – high volume: 2.5±0.8 yrs, low volume: 1.8±0.9 yrs) | Compare the effects of body composition and performance considering distinct volumes of training                              | Body composition (dual-energy X-ray absorptiometry), Fran performance   | ↑ volume of training was associated with ↑ Fran performance<br>↑ volume of training group had ↑ bone mineral density, lean soft mass and ↓ fat mass percentage                                    | Physical characterization                   |
| Carreker et al. [66]  | U.S                       | >6 months                               | N=11 (age: 27.2±3.3 yrs, training experience: 3.9±2.6 yrs)  | Test the physiological predictors of Murph challenge  | Body composition (dual-energy X-ray absorptiometry), upper and lower body strength, body endurance, anaerobic power, $\dot{V}O_{2\text{max}}$ , Murph performance | Body fat percentage was associated with Murph performance ( $r=0.72$ )<br>Run time (during Murph) was associated with relative anaerobic power ( $r=-0.63$ ) and ↓ anaerobic fatigue ( $r=0.64$ ) | Physical characterization                   |
| Cardenosa et al. [67] | Spain                     | >1 yr                                   | N=10 males (age: 30.4±5.4 yrs, training experience: NR)   | Examine the effects of CrossFit® participation (6 weeks) on physical performance, body composition and biochemical parameters | Physical performance, biochemical parameters  | ↔ in physical performance after 6-weeks of CrossFit® training<br>↑ fat oxidation rate (63%)<br>↓ carbohydrate oxidation (27%)<br>↑ lactate dehydrogenase (27%)                                    | Physical and physiological characterization |

**Table 1** (continued)

| Study                      | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics  | Aim  | Main methodologies/ variables extracted   | Main results   | Main topic                     |
|----------------------------|---------------------------|---|---|--|---|--|--------------------------------|
| Timón et al. [68]          | Spain                     | >1 yr                                   | N=12 (age: 30.0±5.4 yrs, training experience: NR)   | Test physical performance and biochemical parameters after two different types of training (e.g., as many rounds as possible, rounds for time) and to examine recovery time 24 and 48 h after these two workouts | Lactate, RPE, HR, two types of workout performances (as many rounds as possible, rounds for time)   | Physiological different between the type of workouts were found: Rounds for time ↑ blood lactate, heart rate, blood glucose compared to as many rounds as possible<br>After workout sessions ↑ hepatic transaminases, CK, blood glucose and ↓ physiological performance<br>After 48 h, values returned to baseline | Compare different workouts     |
| Poderoso et al. [69]       | Brazil                    | >6 months                               | N=17 males, N=12 females (age: 35.3±10.4 yrs, training experience – males: 0.8±0.2 yrs, females: 0.7±0.2 yrs) | Examine the chronic effects (over six months) in CrossFit® practice on hormonal parameters   | Testosterone, cortisol, testosterone:cortisol ratio   | Testosterone ↑ after 6 months and values ↓ among males in comparison to females<br>Cortisol ↓ all time points in comparison to initial levels<br>↓ testosterone:cortisol ratio in females compared to males  | Physiological characterization |
| Martínez-Gómez et al. [70] | Spain                     | 1 yr                                    | N=23 males (age: 33.0±7.0 yrs, training experience: NR)   | Test the association between CrossFit® performance and back squat exercise (i.e. strength and power)   | Sum of scores in specific WODs, peak and mean power (squat), RM squat                               | Robust correlations ( $r=0.47$ o $0.69$ ) were found between squat variables analyzed and CrossFit® performance<br>↑ correlations were noted for absolute and relative RM, peak and mean adjusted for body weight  | Physical characterization      |
| Fetito et al. [71]         | U.S.                      | >2 yrs                                  | N=15 males, N=14 females (age: 28.6±5.7 yrs, training experience: NR)   | Assess the physiological response after Wingate trials with short and active recovery periods  | Four Wingate trials, $\text{VO}_{2\text{max}}$ respiratory exchange ratio, rate of perceived effort | The explanation of 15 min as many rounds as possible ↑ as the participants progress from the first to the third trial of Wingate test  | Physiology – recovery          |

**Table 1** (continued)

| Study                   | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics  | Aim  | Main methodologies/variables extracted  | Main results   | Main topic                     |
|-------------------------|---------------------------|---|---|--|---|--|--------------------------------|
| Dexheimer et al. [72]   | US                        | 1 yr                                    | N=12 males, N=5 females (age: 28.0±5.0 yrs, training experience – males: 4.1±2.7 yrs, females: 2.5±1.5 yrs) | Determine the importance of physiological indicators on CrossFit® performance      | $\text{VO}_{2\text{max}}$ , 3 min maximal test, Wingate test, CrossFit® total, CrossFit® benchmarks | Back squat explained 4.2% of FRAN performance<br>$\text{VO}_{2\text{max}}$ explained 68% of Nancy performance<br>Peak power (derived from Wingate) explained 57% of CrossFit® total performance  | Physiological characterization |
| De Oliveira et al. [73] | Brazil                    | >3 months                               | 10 male and 6 female CrossFit® participants (age: 32. ±6.4 yrs, training experience: 2.4—20 yrs)            | Describe the HR variability among CrossFit® athletes                               | HR variability  | Sympathetic activity was frequent in CrossFit® participants  | Physiological characterization |
| Bartbieri et al. [74]   | Brazil                    | NR                                      | 4 male and 4 female CrossFit® participants (age: 28.6±4.3 yrs, training experience: 1.6±0.7 yrs)            | Compare different types of training (CrossFit®, recreational trained)              | Body composition (skinfolds), cardiorespiratory test  | CrossFit® ↑ lean mass than resistance training<br>CrossFit® ↓ second ventilatory threshold, power of first and second ventilatory threshold than resistance training<br>Heart rate recovery ↓ CrossFit® group than resistance training | Compare different workouts     |
| Tibana et al. [75]      | Brazil                    | >6 months                               | 8 male CrossFit® participants (age: 28.1±5.4 yrs, training experience: 3.8±1.4 yrs)                         | Examine the adequacy of rate of perceived exertion on the control of the intensity | RPE, HR, lactate, number of repetitions   | The RPE and lactate during the all-out sessions were ↑ than the RPE6<br>↔ heart rate area under the curve and RPE6 sessions<br>RPE and lactate were significantly correlated<br>RPE and heart rate were not related                    | Testing                        |

**Table 1** (continued)

| Study                | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics   | Aim   | Main methodologies/variables extracted                             | Main results   | Main topic                                  |
|----------------------|---------------------------|---|--|---|--|--|---|
| Tibana et al. [76]   | Brazil                    | >6 months                               | 9 healthy male CrossFit® participants (age: 27.7±3.2 yrs, training experience: NR)     | Test the acute and effects of CrossFit® sections with different characteristics on physiological parameters | Lactate, RPE, HR   | Shorter sessions ↑ lactate ( $15.9 \pm 2.2 \text{ mmol L}^{-1} \cdot \text{min}^{-1}$ ) compared to longer sessions ( $12.6 \pm 2.6 \text{ mmol L}^{-1} \cdot \text{min}^{-1}$ )<br>Lactate ↑ recovery period for both training sessions in comparison to pre-exercise<br>HR ↔ in both type of sessions<br>RPE was ↔ in both type of sessions after exercise | Physical and physiological characterization |
| Tibana et al. [77]   | Brazil                    | >6 months                               | 30 male CrossFit® participants (age: 27.2±33.0 yrs, training experience: 27.1±4.1 yrs) | Validate the perceived exertion tool during CrossFit® sections  | RPE, HR  | ↑ correlation between Edwards-TRIMP and RPE in all time frames (0, 10, 20 and 30-min post-exercise)  | Testing                                     |
| Tibana et al. [78]   | Brazil                    | >6 months                               | 22 male CrossFit® participants (age: 29.6±4.4 yrs, training experience: 24.0±0.9 yrs)  | Examine the relationship between two three different movements – squat (back, front), snatch and clean      | Strength (back squat, front squat, clean)                          | ↑ snatch, clean ↑ back squat, front squat  | Physical characterization                   |
| Serafini et al. [79] | U.S.                      | Athletic profile in CrossFit® Games     | 3000 male and female participants  | Determine the utility of self-reported variables distinguished the 2016 CrossFit® profile                   | Self-reported performance measures, performance CrossFit@open 2016 | Successful athletes were better in physical condition and skill<br>Lower-level athletes should focus on the development strength and power after attaining a sufficient proficiency of sport-specific skills   | Physical characterization                   |

**Table 1** (continued)

| Study                    | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics  | Aim  | Main methodologies/ variables extracted                | Main results   | Main topic                     |
|--------------------------|---------------------------|---|---|--|--|--|--------------------------------|
| Mangine et al. [80]      | U.S                       | >2 yrs                                  | 5 male CrossFit® participants (age: 34.4±3.8 yrs, training experience: NR)  | Examine metabolic indicators across 5-weeks of CrossFit® international competition | Testosterone, cortisol, testosterone:cortisol ratio    | Testosterone ↑ immediately after training from week 2 to week 5<br>The ↑ on testosterone were also noted 60 min after the session on week 3 and week 5<br>↑ concentrations of cortisol were noted immediately and 30 min after training in all weeks                                 | Physiological characterization |
| Mangine et al. [81]      | U.S                       | Athletic profile in CrossFit® Games     | 133,857 male and female participants  | Develop specific benchmarks for different type of workouts                         | Performance data                                       | Performance benchmarks: Fran male: 250±106 s, Fran female: 331±181 s, Grace male: 180±90 s, Grace female: 213±96 s, Helen male: 9.5±1.9 min, Helen female: 11.1±2.4 min, F50 male: 24.4±5.9 min, F50 female: 27.3±6.9 min, FGB male: 335±65 repetitions, FGB female: 292±62 females) | Physical characterization      |
| Prado Dantas et al. [82] | Brazil                    | > 1 yr                                  | 10 male CrossFit® participants (age: 29.0±6.3 yrs, training experience: NR)   | Examine the effects of CrossFit® sections in hemodynamic indicators                | Hemodynamic variables                                  | Differences were found immediately after the training sessions and 40 min later in systolic blood pressure<br>Diastolic blood pressure was significantly different between groups 40 min after the sessions  | Physiological characterization |
| Ouellette et al. [83]    | U.S                       | >6 months                               | 5 male and 10 female CrossFit® participants (age – male: 30.0±8.0 yrs, female: 32.0±5.0 yrs, training experience: NR) | Determine if upright is appropriate during the recovery process between sets       | Work rate, heart rate, respiratory rate, oxygen uptake | Statistical differences were found between passive and active recovery strategies<br>Passive strategies (i.e. seating) ↑ work rate<br>Passive strategies were associated with ↑ heart rate, respiratory rate and oxygen consumption  | Physical – recovery            |


**Table 1** (continued)

| Study                           | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics  | Aim   | Main methodologies/variables extracted   | Main results  | Main topic                      |
|---------------------------------|---------------------------|---|---|---|--|---|---------------------------------|
| de Sousa et al [84]             | Brazil                    | >1 yr                                   | 13 male CrossFit® participants (age: $26.0 \pm 3.0$ yrs, training experience: NR)                             | Compare physical capacities considering two different types of practice (CrossFit® and recreational participants) | Body size, body composition, fixed bar pull-ups (relative strength of upper limbs), 20-m shuttle run, countermovement jump | Comparison between groups did not show differences on body composition, pull-ups and countermovement jump   | Comparison by competitive level |
| Fernández-Fernández et al. [85] | Spain                     | NR                                      | 10 CrossFit® participants (age: $30.0 \pm 4.2$ yrs, training experience: $1.0 \pm 0.2$ yrs)                   | Examine physiological parameters on different type of WODs (Fran and Cindy)                                       | $\text{VO}_{2\text{max}}$ , HR, blood lactate, energy expenditure and RPE  | Differences on metabolic parameters were found for $\text{VO}_{2\text{max}}$ and energy expenditure in different training sessions  | Compare different workouts      |
| Butcher et al. [86]             | Canada                    | >1 yr                                   | 10 male and 4 female CrossFit® participants (age: $32.7 \pm 5.7$ yrs, training experience: $3.9 \pm 2.0$ yrs) | Determine the impact of physiological and strength measurements on CrossFit® WODs (Grace, Fran, Cindy)            | CrossFit® performance, $\text{VO}_{2\text{max}}$ Wingate test  | Grace and Fran were related with whole body strength (Grace: $r = -0.88$ , Fran: $r = -0.65$ ) and anaerobic threshold (Grace: $r = -0.61$ , Fran: $r = -0.53$ ) No significant associations were noted for Cindy | Physical characterization       |

**Table 1** (continued)

| Study              | Country of contact author | Inclusion criteria (CrossFit® practice) | Sample characteristics  | Aim   | Main methodologies/variables extracted | Main results   | Main topic                     |
|--------------------|---------------------------|---|---|---|--|--|--------------------------------|
| Bellar et al. [10] | U.S.                      | >1 yr                                   | 21 male CrossFit® participants (age: 26.7 ± 4.3 yrs, training experience: NR) | Determine the importance of aerobic and anaerobic capacity on performance | Aerobic capacity, anaerobic power      | Experience participants had ↑ performance in as many rounds as possible and performed ↓ time to complete the workout than unexperienced participants | Physiological characterization |

NR Not reported, HR Heart rate, RPE Rate of perceived exertion, RPD Rate of perceived discomfort, VO<sub>2max</sub> Maximal oxygen uptake, RER Respiratory exchange ratio, RM Maximal repetition, CK Creatine kinase, PRS Perceived recovery status




**Fig. 2** Frequencies of studies published considering the origin of the first author (A) and the athletes' experience (B)

for time workout [45]. Figure 3 (panel B) indicates considerable variation between subjects. Mean heart rate differs substantially when the workouts are compared (Fig. 3, panel C). Higher mean heart rate values were observed in response to Fran's ( $179 \pm 8$  beats·min $^{-1}$ ) and Cindy's ( $182 \pm 7$  beats·min $^{-1}$ ) workouts. The mean value registered for the Murph workout was  $169 \pm 6$  beats·min $^{-1}$ . Comparing the lactate measured immediately after CrossFit® sessions, the highest mean values were observed following rounds for time workouts. The lowest lactate values were noted following Cindy, Fran, and Murph's workouts (Fig. 3, panel D). The values of the ratings of perceived exertion, considering exclusively the studies that used the Borg scale 1–10, showed that rounds for time tend to be classified as the most physically demanding (Fig. 4). Body composition data indicated that male participants (overall mean considering two studies: 10.5%) presented lower values of fat mass percentage than females (overall mean considering two studies: 16.3%) (Fig. 5).


Recovery was assessed following a number of different CrossFit® workouts, with Fran [20, 46], Karen [43], Fight Gone Bad [10], Cindy [41], and Isabel [21] being the most examined workouts, while other studies classified the workouts as many rounds as possible and rounds for time [68]. Recovery was assessed after each workout using measures of physical performance (i.e., jumping, plank time) or physiological outcomes (i.e., heart rate, lactate, CPK). One study investigated the jumping and plank performance until 24 h after the Fran workout [20]. Physiological parameters were measured immediately-, 10-, 20-, and 30-min post-Fran workout [76]. As shown in Fig. 6, 30 min and 24 h after are not sufficient

for the physical and physiological parameters to return to the baseline. The recovery process was also examined following Karen's workout using jumping height and creatine kinase as measures of recovery [43]. Jumping performance was comparable between baseline and 72 h after the Karen workout, whilst creatine kinase was substantially higher 72 h post-Karen workout (baseline:  $151.1 \pm 68.3$ ; 72-h post-training:  $223.0 \pm 86.3$ , Fig. 7). Heart rate and lactate values 30 min after Open 18.4, Fight Gone Bad, and Cindy workouts were also higher than baseline [41, 76]. The same trend was noted in heart rate 5 min after the Isabel workout [21] (Figs. 8, 9, and 10). Although a separate study did not report the differences between the types of workouts, two different types of workouts were used – as many rounds as possible and rounds for time [68]. Physical performance returned to baseline 48 h post-workouts, while the physiological markers were comparable to baseline 72 h after the workout (Figs. 11 and 12).

Table 2 summarizes the studies assessing the chronic effects of CrossFit® participation [67, 69]. Six weeks was not sufficient to improve the physical performance outputs, but 8 weeks of CrossFit® participation resulted in increases in testosterone and decreases in cortisol levels. Using the same participants, only one interventional study assessed the change in physiological response (blood lactate, heart rate, ratings of perceived exertion, and ratings of discomfort) to different types of training (Cindy workout vs. continuous running) [23]. The mean heart rate was comparable in both groups. However, the maximal heart rate, blood lactate, ratings of perceived exertion, and ratings of perceived discomfort were higher during Cindy's workout. Separate investigations



**Fig. 3** Physiological indicators reported according to type of CrossFit® challenge




**Fig. 4** Rate of perceived exertion considering different types of CrossFit® challenges

compared CrossFit® athletes with other groups, including runners [22], sedentary populations [25], athletes completing cross-training exercise [54], physically active cohorts [25], and resistance training participants [12, 84]

in relation to changes in body composition, physiological responses, and physical performance.

Table 3 presents data for body composition and VO<sub>2max</sub>. CrossFit® participants had lower values of fat



**Fig. 5** Mean values of fat mass percentage splitting by sex

mass percentage than sedentary [25], cross-training [54], or physically active individuals [65]. The  $\text{VO}_{2\text{max}}$  of CrossFit® athletes was higher than the sedentary group [25] and lower than the runners [22]. CrossFit® participants also demonstrated superior physical performance (jumping performance, box-jump, pull-ups, push-ups, burpees, maximum speed) than cross-training [54] and resistance training participants [84].

Interventional [27, 34, 62] and observational studies [10, 28, 60] examined variation by competitive level. Two studies compared the mean and maximal heart rate in experienced and less experienced participants after implementing different training sessions [27] and Cindy workouts [62]. Mean values were equivalent between groups. Cognition variables were measured in 32 CrossFit® athletes classified as elite, advanced, and beginner participants after Fran workouts [34]. All groups differed significantly between pre- and post-Fran workout. Additionally, two observational studies showed that CrossFit® participants of elite or advanced levels had lower fat mass values than those at the middle or recreational level [28, 60]. Strength variables were also affected by the competitive level of participants, with more experienced athletes showing higher values in the maximal repetition test [60], isometric mid-thigh pull assessment [28], rate of force development [28], and power [10].

#### Meta-correlation

#### Challenges focused on completing as many rounds as possible

Studies that used specific protocols of CrossFit® (back squat, front squat, snatch, clean and jerk) found a positive





and moderate relationship between challenges focused on completing as many rounds as possible ( $r=0.33$ ; 95% CI: -0.09 to 0.65), which means that participants who lift more weight on strength protocols complete more rounds on the workouts. The magnitude of correlation increased when the specific protocols were expressed per kilogram of body mass ( $r=0.38$ ; 95% CI: 0.03 to 0.65;  $p=0.03$ ) (Figs. 13 and 14). In contrast, non-specific protocols were not associated with performance in as many rounds as possible workout. The values of heterogeneity were high and moderate for specific ( $I^2=77.1\%$ ) and specific protocols relativized for body mass ( $I^2=67\%$ ), respectively.

#### Challenges focused on completing the workout fasting as possible

When the workouts focused on completing the challenge as fast as possible, time and specific CrossFit® protocols were not associated, as shown in Fig. 15. The overall correlation coefficient was positive ( $r=-0.18$ ; 95% CI: -0.36 to 0.01;  $p=0.07$ ), demonstrating that athletes who were faster on challenge performed less strength on specific protocols. Similar results were obtained for non-specific CrossFit® protocols.

#### Specific workouts

Studies with correlation data between protocols and performance were available for the following workouts: Cindy, Fran, Donkey Kong, Grace, Murph, and Nancy. For Murph and Nancy workouts, data was limited, and for this reason, a meta-correlation was not conducted.

**A.****B.****C.****D.**


**Fig. 6** Physical and physiological variation before and after Fran challenge

For the Cindy workout (which comprises completing as many rounds as possible in 20 min of five pull-ups, ten push-ups, and fifteen air squats), only one study (Butcher et al., 2015) tested the association between the number of rounds complete and non-specific (Wingate test and  $\text{VO}_{2\text{max}}$ ) and specific protocols (CrossFit®). As shown in Supplementary Material 1, the combination of different protocols resulted in a non-significant association with rounds performed on Cindy workout ( $r=0.14$ ; 95% CI: -0.15 to 0.41;  $p=0.33$ ).


The Fran challenge involves completing as fast as possible three rounds of 21, 15, and 9 repetitions of two exercises: thrusters and pull-ups. The meta-correlation of

sub-groups (specific and non-specific) noted significant magnitudes of associations between types of protocol and performance (Fig. 16). A negative correlation coefficient indicated that more time to complete the challenge was associated with better performance in protocols. A small and non-significant association between Fran performance and non-specific protocols ( $r=0.24$ ,  $p=0.19$ ) was found. In opposition, specific protocols were moderately associated with performance in the Fran challenge ( $r=-0.44$ ; 95% CI: 0.22 to 0.54;  $p<0.05$ ). The value of heterogeneity was high ( $I^2=75\%$ ).

On the Donkey Kong challenge, CrossFit® participants should complete as fast as possible three rounds




**Fig. 7** Physical and physiological variation before and after Karen challenge




**Fig. 8** Physical and physiological variation before and after Open 18.4 challenge

of 21, 15, and 9 repetitions of burpees, kettlebell swings, and box jumps. After each exercise, participants need to perform six lunges. A negative correlation coefficient indicated that less time to complete the challenge was associated with a better performance in Donkey Kong challenge. Only one study examined the relationship between different protocols and performance in this challenge (Gomez-Landero et al., 2020). For

non-specific protocols, the study included four different protocols: sit-ups, hand dynamometers,  $\text{VO}_{2\text{max}}$  estimated from a shuttle-run test, and peak power derived from countermovement jumps. Specific protocols used were pull-ups, bench presses, and squats. Non-specific protocols were significantly related with performance, while the overall magnitude of correlation in specific protocols was small and non-significant ( $r = -0.26$ ; 95% CI: -0.06 to 0.53;  $p = 0.11$ ) (Supplementary Material 2).



**Fig. 9** Physiological variation before and after Fight Gone Bad challenge




**Fig. 10** Physiological variation before and after Cindy challenge


The Grace challenge consists of performing 30 repetitions of clean and jerk as fast as possible, which means a negative correlation represents less time to complete the workout. Non-specific protocols were not associated with performance on Grace performance. On the other hand, total strength, strict press, deadlift, and back squat exercises were related to Grace performance ( $r=0.478$ ; 95% CI: 0.177 to 0.697;  $p=0.003$ ). The heterogeneity across studies was moderate ( $I^2=62\%$ ) (Fig. 17).

## Discussion

The aim of this scoping review was to characterize the physical demands and physiological responses to CrossFit®. The findings and potential gaps in the scientific literature that emerged from the current review were as follows: (1) CrossFit® studies have mainly been undertaken in North and South America; (2) the definition of a CrossFit® athlete is not clear in the literature with different cut-off values being used to include participants in the studies; (3) a limited number of studies focused on



**Fig. 11** Physical and physiological variation before and after as many rounds as possible workouts



**Fig. 12** Physical and physiological variation before and after rounds for time workouts

**Table 2** Studies<sup>a</sup> that described the chronic effects of CrossFit® participation

| Study                 | Time of practice | Output                  | Result |
|-----------------------|------------------|-------------------------|--------|
| Poderoso et al. [69]  | 2 months         | Testosterone            | ↑      |
|                       | 4 months         |                         | ↑      |
|                       | 6 months         |                         | ↑      |
|                       | 2 months         | Cortisol                | ↓      |
|                       | 4 months         |                         | ↓      |
|                       | 6 months         |                         | ↓      |
| Cardeñosa et al. [67] | 6 weeks          | Heart rate <sup>b</sup> | ↔      |
|                       |                  | Maximal power           | ↓      |
|                       |                  | VO <sub>2max</sub>      | ↔      |
|                       |                  | Sum of six skinfolds    | ↑      |

<sup>a</sup> Data from da Silveira Castanheira et al. [18] was not provided. <sup>b</sup> heart rate was derived from an incremental test; ↔ = no difference, ↑ = increased, ↓ = reduced

characterizing the physiological and physical parameters of different workouts; (4) body composition data suggest that males have less fat mass percentage than females; (5) recovery strategies for CrossFit® should be investigated


in order to optimize weekly performance and physiological markers; (6) the literature relation to the chronic effects of CrossFit® is scarce, although the study that did exist in this area demonstrated that six weeks was not sufficient to promote significant changes in physical and physiological parameters, while eight weeks led to increases in testosterone and decrements in cortisol; (7) in comparison to other exercise modalities (i.e., resistance training, endurance), CrossFit® elicits greater benefits to body composition and maximal oxygen uptake; (8) it was not possible to determine a unique predictor of CrossFit® performance; (9) movements specific to CrossFit® seem to be more related to CrossFit® performance than non-specific protocols.

An early study compared three groups across ten weeks, with participants either undertaking aerobic training ( $n=8$ ), resistance training ( $n=8$ ), or concurrent training ( $n=7$ ) [88]. Concurrent training involves the inclusion of resistance training (to gain strength, hypertrophy, and power) combined with aerobic exercise (to enhance endurance) [89]. A reduction in lower body strength was found in the concurrent training group

**Table 3** Observational studies comparing Crossfit® with other activities or control group


| Variable   | Study                | Crossfit® vs. comparator | Crossfit®   | Comparator  |
|--|----------------------|--------------------------|-------------|-------------|
|  |                      |                          | Mean ± sd   | Mean ± sd   |
| Fat mass, %  |                      |                          |             |             |
|  | Carvalho et al. [22] | vs. runners              | 19.0 ± 6.3  | 18.2 ± 6.3  |
|  | Pearson et al. [25]  | vs. sedentary            | 18.6 ± 3.8  | 30.3 ± 8.4  |
|  | Fernando et al. [54] | vs. crosstraining        | 16.5 ± 27.5 | 18.0 ± 3.0  |
|  | Cavedon et al. [65]  | vs. physical active      | 12.9 ± 2.7  | 18.9 ± 3.3  |
|  | Barbieri et al. [74] | vs. resistance training  | 11.8 ± 4.9  | 14.1 ± 5.3  |
|  | de Sousa et al. [84] | vs. resistance training  | 13.6 ± 4.6  | 11.9 ± 4.1  |
| Lean mass, kg  |                      |                          |             |             |
|  | Pearson et al. [25]  | vs. sedentary            | 64.6 ± 11.7 | 50.7 ± 6.3  |
|  | Cavedon et al. [65]  | vs. physical active      | 65.0 ± 6.4  | 64.8 ± 6.2  |
|  | Barbieri et al. [74] | vs. resistance training  | 59.8 ± 10.7 | 49.6 ± 8.1  |
| VO <sub>2max</sub> , ml kg <sup>-1</sup> min <sup>-1</sup> |                      |                          |             |             |
|  | Carvalho et al. [22] | vs. runners              | 41.8 ± 5.4  | 50.0 ± 10.8 |
|  | Pearson et al. [25]  | vs. sedentary            | 43.3 ± 3.9  | 31.1 ± 3.8  |
|  | Barbieri et al. [74] | vs. resistance training  | 42.5 ± 5.3  | 44.4 ± 5.5  |
|  | de Sousa et al. [84] | vs. resistance training  | 52.5 ± 5.6  | 46.0 ± 5.8  |

sd Standard deviation

**Fig. 13** Meta-correlation between specific protocols and CrossFit® workouts classified as many rounds as possible. F (female); M (male). Note: positive indicates positive performance

compared to resistance exercise. It was hypothesized that aerobic training negatively impacted the resistance training adaptations, termed the “interference effect” [66]. In a meta-analysis that combined 21 studies, resistance training promoted higher gains in hypertrophy, strength, and power than concurrent protocols [90]. This study also concluded that the type and volume of endurance training impact the “interference effects” of resistance training [90]. While in the meta-analysis, the details of studies

were not presented, the participants of the concurrent training group included in the original research about the “interference effect” completed the resistance training and endurance protocol separated by two hours [88]. The training sessions of CrossFit® incorporated both endurance and resistance exercises within the same session, which suggests the “interference effect” cannot be generalized for CrossFit® participants. Nevertheless, more recent studies about the short-term effects of concurrent




**Fig. 14** Meta-correlation between specific protocols normalized for body weight and CrossFit® workouts classified as many rounds as possible. F (female); M (male). Note: positive indicates positive performance

training on muscle hypertrophy showed contradictory findings questioning the “interference effect” theory [87, 91, 92]. A parallel study compared two different conditions (resistance training in isolation vs. concurrent training [i.e., combining cycling activities with resistance training]) across a 7-week program on muscle size and specific indicators of protein synthesis and degradation [91]. Muscle fiber area increased significantly in the concurrent training group, whilst negligible changes were noted in the resistance training group. The levels of the mechanistic target of rapamycin (i.e., an indicator of muscle mass development) were also raised in concurrent training, highlighting the anabolic effects when endurance and resistance activities were combined [91]. In the present review, CrossFit® participants had higher mean values of lean mass in comparison to physically active [65] and resistance training [74] participants, suggesting the combination of resistance training and endurance exercise within the same training session could potentiate the development of muscle mass. Participation in CrossFit® sessions over 8 weeks also demonstrated increases in testosterone and decreases in cortisol.

The main findings pertaining to concurrent training are modulated by the training status of participants and the methodologies used to assess changes in muscle strength and hypertrophy [5]. Three to nine months were proposed to classify an athlete as “trained” for a resistance and endurance athlete [5]. Although most of the studies included in the present review attained these criteria, the literature focused exclusively on the best athletes was scarce. Strength, power, and body composition distinguished elite athletes from lower competitive levels [10,

28, 60]. Moreover, considering the CrossFit® Open allows everyone to participate, significant variability in physical performance is expected. Future studies should focus on examining participants considering the different phases of CrossFit® competition: CrossFit® Open, quarterfinals, semi-finals, and CrossFit® Games. It might then be possible to discriminate participants according to training status and not focus exclusively on training time. Specificity was another concept claimed to define training status, particularly task-specific activities related to maximal strength. Hypertrophy is not exercise-dependent [5, 93], while changes in strength are exercise- and intensity-dependent (i.e., specific).

The relationship between CrossFit® performance and different protocols seems specific, with movements often carried out during CrossFit® sessions (e.g., back squat, front, deadlift, clean, clean and jerk) being more related to performance. In contrast, non-specific protocols were rarely associated with CrossFit® performance. For those who train CrossFit® athletes, the application of specific protocols has more relevance. Whereas, on the other hand, the results of the current meta-correlation also showed that it is difficult to generalize a particular test for all types of workouts. This is in line with a systematic review that described back squat and total body strength as the main variables to explain performance. However, variation in the results across the 21 studies included indicates that a consensus about predictors could not be generalized [9]. Considering the variability of CrossFit® in terms of exercises and intensity, this point is not surprising. The variability of physical and physiological indicators to explain the performance in CrossFit® workouts




**Fig. 15** Fig. 14. Meta-correlation between specific protocols and CrossFit® workouts classified as time to complete. F (female); M (male). Note: positive indicates positive performance

was noted, which indicates that the predictors of a typical endurance workout should not be generalized for a resistance workout [21, 28, 42, 66, 79]. In order to support coaches in the monitoring, quantification, and regulation of training load, future studies need to investigate the physical and physiological characteristics of other specific workouts.


A considerable number of studies used physical and physiological outputs to examine recovery after a CrossFit® workout [20, 41, 43, 68, 88]. In general, 48–72 h

were insufficient to obtain the baseline values of physical and physiological markers, and these studies only focused on a specific challenge. A typical CrossFit® session includes other components rather than the workout of the day (e.g., strength, mobility, stability, skill). In order to prevent fatigue and optimize performance, research is needed on which recovery strategies for CrossFit® athletes are needed [6, 14].

The inclusion of papers solely written in English, Portuguese, and Spanish is a limitation of the current



**Fig. 16** Meta-correlation between specific, non-specific protocols and Fran performance. Note: positive indicates positive performance



**Fig. 17** Meta-correlation between specific, non-specific protocols and Grace performance. Note: positive indicates positive performance

review. Studies about injuries or psychological variables were not considered since these topics were previously discussed in the literature [13, 94]. The non-uniform criteria relating to training experience required to qualify as a CrossFit® athlete is the major limitation of the eligible studies. Consequently, the participants' level must be discriminated in the sampling description. In most of the studies, a specific workout's physical or physiological description did not always consider the entire training session. Consequently, future studies should investigate training sessions' physical and

physiological aspects individually or combined in microcycles or mesocycles.

## Conclusion

CrossFit® seems to align with the recent benefits described in concurrent training, although they are modulated by training status and specificity of exercise. The definition of CrossFit® athlete needs to be considered in future studies since everyone can perform CrossFit Open®. In this competition, significant variability in performance and participants' characteristics are observed,

influencing the interpretation of results. The correct manipulation of the training load is an additional issue for coaches in order to optimize performance and prevent fatigue. Coaches should be aware that the design and implementation of CrossFit® programs require specific information about the metabolic demands of each workout. Consequently, they should use training tools to control the volume and intensity of training to manage the training load. In order to interpret performance, protocols with specific CrossFit® movements should be routinely applied, even though it was challenging to obtain a test for all workouts. Therefore, further research needs to be conducted to characterize workouts that induce distinct physical and physiological responses.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13102-024-00986-3>.

Supplementary Material 1.

Supplementary Material 2.

## Acknowledgements

Not applicable.

## Authors' contributions

DS, DVM, ERG, HS conceptualized the manuscript. DS, DVM, ERG, HS wrote the protocol and methodology. DVM, AR, CA, DSB, NS used and managed the software. DVM, ERG, AS, AF, HS organized the data. DVM, AR, CA, NS, AS writing the original draft. DS, DVM, CM, AF, ERG, HS wrote and reviewed the manuscript.

## Funding

No sources of funding were used to assist in the preparation of this article.

## Availability of data and materials

All data generated or analysed during this study are included in this published article and its supplementary information file.

## Declarations

### Ethics approval and consent to participate

Not applicable.

### Consent for publication

Not applicable.

### Competing interests

The authors declare no competing interests.

Received: 22 July 2024 Accepted: 10 September 2024

Published online: 20 September 2024

## References

- Glassman G. Understanding CrossFit CrossFit J. 2007;56:1–2.
- Schlegel P. CrossFit® Training Strategies from the Perspective of Concurrent Training: A Systematic Review. J Sports Sci Med. 2020;19(4):670–80.
- Glassman G. What is fitness CrossFit J. 2002;3:1–11.
- Glassman G. Benchmark workouts CrossFit J. 2003;13:1–5.
- Fyfe JJ, Loenneke JP. Interpreting Adaptation to Concurrent Compared with Single-Mode Exercise Training: Some Methodological Considerations. Sports Med. 2018;48(2):289–97.
- Halson SL. Monitoring training load to understand fatigue in athletes. Sports Med. 2014;44 Suppl 2(Suppl 2):S139–47.
- About the Games. 2023. [<https://games.crossfit.com/about-the-games>].
- Edmonds W. Is the CrossFit Open the biggest sporting competition on Earth? CNN Sports. 2018.
- Meier N, Schlie J, Schmidt A. CrossFit®: “Unknowable” or Predictable?—A Systematic Review on Predictors of CrossFit® Performance. Sports (Basel). 2023;11(6):112.
- Bellar D, Hatchett A, Judge LW, Breaux ME, Marcus L. The relationship of aerobic capacity, anaerobic peak power and experience to performance in CrossFit exercise. Biol Sport. 2015;32(4):315–20.
- Meyer J, Morrison J, Zuniga J. The Benefits and Risks of CrossFit: A Systematic Review. Workplace Health Saf. 2017;65(12):612–8.
- Ángel Rodríguez M, García-Calleja P, Terrados N, Crespo I, Del Valle M, Olmedillas H. Injury in CrossFit®: A Systematic Review of Epidemiology and Risk Factors. Phys Sportsmed. 2022;50(1):3–10.
- Dominski FH, Serafim TT, Siqueira TC, Andrade A. Psychological variables of CrossFit participants: a systematic review. Sport Sci Health. 2021;17(1):21–41.
- de Souza RAS, da Silva AG, de Souza MF, Roschel H, Silva SF, Saunders B. A Systematic Review of CrossFit® Workouts and Dietary and Supplementation Interventions to Guide Nutritional Strategies and Future Research in CrossFit®. Int J Sport Nutr Exerc Metab. 2021;31(2):187–205.
- Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). Cochrane handbook for systematic reviews of interventions. 2nd Edition. Chichester: Wiley; 2019.
- Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ. 2021;372: n71.
- Wojtyniak JG, Britz H, Selzer D, Schwab M, Lehr T. Data Digitizing: Accurate and Precise Data Extraction for Quantitative Systems Pharmacology and Physiologically-Based Pharmacokinetic Modeling. CPT Pharmacometrics Syst Pharmacol. 2020;9(6):322–31.
- Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies in sports medicine and exercise science. Med Sci Sports Exerc. 2009;41(1):3–13.
- Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. Stat Med. 2002;21(11):1539–58.
- Rios M, Becker KM, Monteiro AS, Fonseca P, Pyne DB, Reis VM, Moreira-Gonçalves D, Fernandes R. Effect of the Fran CrossFit Workout on Oxygen Uptake Kinetics, Energetics, and Postexercise Muscle Function in Trained CrossFitters. Int J Sports Physiol Perform. 2024;19(3):299–306.
- Rios M, Becker KM, Cardoso F, Pyne DB, Reis VM, Moreira-Gonçalves D, Fernandes RJ. Assessment of Cardiorespiratory and Metabolic Contributions in an Extreme Intensity CrossFit® Benchmark Workout. Sensors (Basel). 2024;24(2):513.
- Carvalho LL, Costa DA, N, Mansour KM, Simonis JG, Teixeira L, Gonçalves DP, Rekziegel MB, Possuelo LG, Valim ARDE. Effects of CrossFit® and street running practice on anthropometric, lipids parameters, cardiorespiratory fitness and sleep quality. J Sports Med Phys Fitness. 2024;64(1):1–6.
- Santos DAT, Morais NS, Viana RB, Costa GCT, Andrade MS, Vancini RL, Weiss K, Knechtel B, de Lira CAB. Comparison of physiological and psychobiological acute responses between high intensity functional training and high intensity continuous training. Sports MedHealth Sci. 2023. <https://doi.org/10.1016/j.smhs.2023.10.006>.
- Rios M, Zaccaria R, Azevedo R, Fonseca P, Pyne DB, Reis VM, Moreira-Gonçalves D, Fernandes RJ. Bioenergetic Analysis and Fatigue Assessment During the Fran Workout in Experienced Crossfitters. Int J Sports Physiol Perform. 2023;18(7):786–92.
- Pearson RC, Olenick AA, Jenkins NT. Metabolic response during high-intensity interval exercise and resting vascular and mitochondrial function in CrossFit participants. Kinesiology. 2023;55(2):228–44.
- Párraga-Montilla JA, Cabrera Linares JC, Jiménez Reyes P, Moyano López M, Serrano Huete V, Morcillo Losa JA, Latorre Román PA. Force–velocity profiles in CrossFit athletes: A cross-sectional study considering sex, age, and training frequency. Balt J Health Phys Act. 2023;15(1):5.
- Meier N, Sietmann D, Schmidt A. Comparison of cardiovascular parameters and internal training load of different 1-h training sessions in non-elite CrossFit® athletes. J Sci Sport Exerc. 2023;5(2):130–41.

28. Manrique JEH, Chavarria WEBM, Sánchez WGV, Velásquez CAA. Are there differences in maximal strength, flexibility, and body composition in CrossFit® competitors according to their category? *Retos.* 2023;47:866–75.
29. Mangine GT, Zeitz EK, Dexheimer JD, Hines A, Lively B, Kliszczewicz BM. Pacing Strategies Differ by Sex and Rank in 2020 CrossFit® Open Tests. *Sports (Basel).* 2023;11(10):199.
30. Mangine GT, Grundlingh N, Feito Y. Normative Scores for CrossFit® Open Workouts: 2011–2022. *Sports (Basel).* 2023;11(2):24.
31. Mangine GT, Grundlingh N, Feito Y. Differential improvements between men and women in repeated CrossFit open workouts. *PLoS ONE.* 2023;18(11):e0283910.
32. Linhares M, Façanha C, Teixeira M, Alves KS, Pires T, Coswig V, Cabido C, Fermino RC, Oliveira S, de Faria Souza TM, Aidar FJ, Acioli T, Cirilo-Sousa MS, da Rabello Lima LC, Bertú F, Assumpção CO, Banja T. Examining strength, muscular power, and maximal performance in the power clean among CrossFit® practitioners. *J Phys Educ Sport.* 2023;23(11):3119–26.
33. da Silveira Castanheira LF, Penna EM, Franco ECS, Coswig VS. Training load monitoring and physiological responses to RX CrossFit® training. *J Phys Educ Sport.* 2023;23(5):1076–85.
34. Brito MA, Fernandes JR, De Carvalho PHB, Brito CJ, Aedo-Muñoz E, Soto DAS, Miarka B. Acute Effect of a Cross-Training Benchmark on Psycho-physiological Factors of Cross-Training According to Performance. *Sport Mont.* 2023;21(2):9–15.
35. Barreto AC, Medeiros AP, da Silva AG, de Souza Vale RG, Vianna JM, Alkimir R, Serra R, Leitão L, Reis VM, da Silva Novaes J. Heart rate variability and blood pressure during and after three CrossFit® sessions. *Retos.* 2023;47:311–6.
36. Schlegel P, Krehký A. Performance Sex Differences in CrossFit®. *Sports (Basel).* 2022;10(11):165.
37. Menargues-Ramírez R, Sospedra I, Holway F, Hurtado-Sánchez JA, Martínez-Sanz JM. Evaluation of Body Composition in CrossFit® Athletes and the Relation with Their Results in Official Training. *Int J Environ Res Public Health.* 2022;19(17):11003.
38. Martínez-Gómez R, Valenzuela PL, Lucia A, Barranco-Gil D. Comparison of Different Recovery Strategies After High-Intensity Functional Training: A Crossover Randomized Controlled Trial. *Front Physiol.* 2022;13: 819588.
39. Mangine GT, McDougle JM, Feito Y. Relationships Between Body Composition and Performance in the High-Intensity Functional Training Workout "Fran" are Modulated by Competition Class and Percentile Rank. *Front Physiol.* 2022;13: 893771.
40. Mangine GT, Seay TR. Quantifying CrossFit®: Potential solutions for monitoring multimodal workloads and identifying training targets. *Front Sports Act Living.* 2022;4: 949429.
41. Forte LDM, Freire YGC, Júnior JS, Melo DA, Meireles CLS. Physiological responses after two different CrossFit workouts. *Biol Sport.* 2022;39(2):231–6.
42. Dias MR, Vieira JG, Pissolato JC, Heinrich KM, Vianna JM. Training load through heart rate and perceived exertion during CrossFit®. *Rev Bras Med Esporte.* 2022;28:315–9.
43. Sousa Neto IVD, Sousa NMFD, Neto FR, Falk Neto JH, Tibana RA. Time course of recovery following CrossFit® Karen Benchmark workout in trained men. *Front Physiol.* 2022;2022(13): 899652.
44. Conde TF, Silva MRDS, Caobianco J, Robalino J, Ferreira JC. Sensitivity of operational tests to training load in CrossFit®. *J Phys Educ Sport.* 2022;22(6):1493–8.
45. Toledo R, Dias MR, Toledo R, Ertides R, Pinto DS, Reis VM, Novaes JS, Vianna JM, Heinrich KM. Comparison of Physiological Responses and Training Load between Different CrossFit® Workouts with Equalized Volume in Men and Women. *Life (Basel).* 2021;11(6):586.
46. Tibana RA, de Sousa Neto IV, Sousa NM, Romeiro C, Hanai A, Brandão H, Dominski FH, Voltarelli FA. Local Muscle Endurance and Strength Had Strong Relationship with CrossFit® Open 2020 in Amateur Athletes. *Sports (Basel).* 2021;9(7):98.
47. Schlegel P, Režný L, Fialová D. Pilot study: Performance-ranking relationship analysis in Czech crossfitters. *J Hum Sport Exerc.* 2021;16(1):187–98.
48. Ponce-García T, Benítez-Porres J, García-Romero JC, Castillo-Domínguez A, Alveró-Cruz JR. The Anaerobic Power Assessment in CrossFit® Athletes: An Agreement Study. *Int J Environ Res Public Health.* 2021;18(16):8878.
49. Peña J, Moreno-Doutres D, Peña I, Chulvi-Medrano I, Ortegón A, Aguilera-Castells J, Buscà B. Predicting the Unknown and the Unknowable. Are Anthropometric Measures and Fitness Profile Associated with the Outcome of a Simulated CrossFit® Competition? *Int J Environ Res Public Health.* 2021;18(7):3692.
50. Mota MR, Brandao HCP, Alencastro G, Elias R, Ribeiro A, de Araujo Ribeiro AL, Chaves SN, Cleto F, Silva AS, Clael S. Glycemia Analysis in Two Different CrossFit [R] Benchmark Protocols. *J Exerc Physiol Online.* 2021;24(2):1–9.
51. Meier N, Rabel S, Schmidt A. Determination of a CrossFit® Benchmark Performance Profile. *Sports (Basel).* 2021;9(6):80.
52. Mangine GT, Feito Y, Tankersley JE, McDougle JM, Kliszczewicz BM. Workout Pacing Predictors of CrossFit® Open Performance: A Pilot Study. *J Hum Kinet.* 2021;78:89–100.
53. Leitão L, Dias M, Campos Y, Vieira JG, San't Ana L, Telles LG, Tavares C, Maziní M, Noyes J, Vianna J. Physical and Physiological Predictors of FRAN CrossFit® WOD Athlete's Performance. *Int J Environ Res Public Health.* 2021;18(8):4070.
54. Fernando W, Santos W, Barbieri J, de Medeiros Lima LE, Miguel H, Guedes D Jr, Silva RP, Marchioni E, Moriggi JRR. Physical capacities and anthropometric measures between crossfit® practitioners and cross-training. *Multidiscip Res J.* 2021;3(2): e0201006.
55. Fernández-Lázaro D, Mielgo-Ayuso J, Novo DFZS, Lázaro-Asensio MP, Sánchez-Serrano N, Fernández-Lázaro CI. Athletic, muscular and hormonal evaluation in CrossFit® athletes using the Elevation Training Mask. *Arch Med Deporte.* 2021;38:274–81.
56. Bustos-Vivencias BJ, Luna LAD, Osorio RDM, Parra AJO, Acevedo-Mindiola AA, García Yerena CE. High-intensity functional training: Association of body fat with cardiorespiratory fitness. *Revista Cubana de Medicina Militar.* 2021;50(2):1–13.
57. Pritchard HJ, Keogh JW, Winwood PW. Tapering practices of elite CrossFit athletes. *Int J Sports Sci Coach.* 2020;15(5–6):753–61.
58. Martínez-Gómez R, Valenzuela PL, Alejo LB, Gil-Cabrera J, Montalvo-Pérez A, Talavera E, Lucia A, Moral-González S, Barranco-Gil D. Physiological Predictors of Competition Performance in CrossFit Athletes. *Int J Environ Res Public Health.* 2020;17(10):3699.
59. Mangine GT, Tankersley JE, McDougle JM, Velazquez N, Roberts MD, Esmat TAM. Predictors of CrossFit Open Performance. *Sports (Basel).* 2020;8(7):102.
60. Mangine GT, Stratton MT, Almeda CG, Roberts MD, Esmat TA, VanDusseldorp T, Feito Y. Physiological differences between advanced CrossFit athletes, recreational CrossFit participants, and physically-active adults. *PLoS ONE.* 2020;15(4): e0223548.
61. Gómez-Landero LA, Frías-Menacho JM. Analysis of Morphofunctional Variables Associated with Performance in CrossFit® Competitors. *J Hum Kinet.* 2020;73:83–91.
62. Gomes JH, Mendes RR, Franca CS, Silva-Grigoletto MES, Silva DRP, Antonioli AR, Oliveira E, Silva AM, Quintans-Júnior L. Acute leucocyte, muscle damage, and stress marker responses to high-intensity functional training. *PLoS One.* 2020;15(12): e0243276.
63. Faelli E, Bisio A, Codella R, Ferrando V, Perasso M, Panascì M, Saverino D, Ruggeri P. Acute and Chronic Catabolic Responses to CrossFit® and Resistance Training in Young Males. *Int J Environ Res Public Health.* 2020;17(19):7172.
64. Dexheimer JD, Schroeder ET, Sawyer BJ, Pettitt RW, Torrence WA. Total Body Strength Predicts Workout Performance in a Competitive Fitness Weightlifting Workout. *J Exerc Physiol Online.* 2020;23(4):95–104.
65. Cavedon V, Milanese C, Marchi A, Zancanaro C. Different amount of training affects body composition and performance in High-Intensity Functional Training participants. *PLoS ONE.* 2020;15(8): e0237887.
66. Carreker JD, Grosicki GJ. Physiological Predictors of Performance on the CrossFit "Murph" Challenge. *Sports (Basel).* 2020;8(7):92.
67. Cardeñosa AC, Andrada RT, Cardeñosa MC, Flores SG, Camacho GJO, Serrano MM. Six-months CrossFit training improves metabolic efficiency in young trained men. *Culto Ciência Esporte.* 2020;15(45):421–7.
68. Timón R, Olcina G, Camacho-Cardeñosa M, Camacho-Cardenosa A, Martínez-Guardado I, Marcos-Serrano M. 48-hour recovery of biochemical parameters and physical performance after two modalities of CrossFit workouts. *Biol Sport.* 2019;36(3):283–9.
69. Poderoso R, Cirilo-Sousa M, Júnior A, et al. Gender Differences in Chronic Hormonal and Immunological Responses to CrossFit®. *Int J Environ Res Public Health.* 2019;16(14):2577.

70. Martínez-Gómez R, Valenzuela PL, Barranco-Gil D, Moral-González S, García-González A, Lucia A. Full-Squat as a Determinant of Performance in CrossFit. *Int J Sports Med.* 2019;40(9):592–6.
71. Feito Y, Giardina MJ, Butcher S, Mangine GT. Repeated anaerobic tests predict performance among a group of advanced CrossFit-trained athletes. *Appl Physiol Nutr Metab.* 2019;44(7):727–35.
72. Dexheimer JD, Schroeder ET, Sawyer BJ, Pettitt RW, Aguinaldo AL, Torrence WA. Physiological Performance Measures as Indicators of CrossFit® Performance. *Sports (Basel).* 2019;7(4):93.
73. de Oliveira FTO, Ramos ACC, Almeida CN, Dos Santos CPC, Oliveira IAA, Mendel MDR, Santos ECL, Dias CMCC. Modulation of heart rate variability in CrossFit® practitioners. *Revista Pesquisa em Fisioterapia.* 2019;9(3):353–60.
74. Barbieri JF, Figueiredo GTDC, Castano LAA, Guimaraes PDS, Ferreira RR, Ahmadi S, Gaspari AF, De Moraes AC. A comparison of cardiorespiratory responses between CrossFit® practitioners and recreationally trained individual. *J Phys Educ Sport.* 2019;19(3):1606–11.
75. Alsamir Tibana R, de Manuel Fraide Sousa N, Prestes J, Nascimento DC, Ernesto C, Neto JHF, Kennedy MD, Voltarelli FA. Is Perceived Exertion a Useful Indicator of the Metabolic and Cardiovascular Responses to a Metabolic Conditioning Session of Functional Fitness? *Sports (Basel).* 2019;7(7):161.
76. Tibana RA, De Sousa NMF, Prestes J, Voltarelli FA. Lactate, Heart Rate and Rating of Perceived Exertion Responses to Shorter and Longer Duration CrossFit® Training Sessions. *J Funct Morphol Kinesiol.* 2018;3(4):60.
77. Tibana RA, de Sousa NMF, Cunha GV, Prestes J, Fett C, Gabbert TJ, Voltarelli FA. Validity of Session Rating Perceived Exertion Method for Quantifying Internal Training Load during High-Intensity Functional Training. *Sports (Basel).* 2018;6(3):68.
78. Tibana RA, Farias DL, Nascimento DC, Silva-Grigoletto ME, Prestes J. Correlation of muscle strength with weightlifting performance in CrossFit® practitioners. *Rev Andaluza Med Deporte.* 2016;11(2):84–8.
79. Serafini PR, Feito Y, Mangine GT. Self-reported Measures of Strength and Sport-Specific Skills Distinguish Ranking in an International Online Fitness Competition. *J Strength Cond Res.* 2018;32(12):3474–84.
80. Mangine GT, Van Dusseldorp TA, Feito Y, Holmes AJ, Serafini PR, Box AG, Gonzalez AM. Testosterone and Cortisol Responses to Five High-Intensity Functional Training Competition Workouts in Recreationally Active Adults. *Sports (Basel).* 2018;6(3):62.
81. Mangine GT, Cebulla B, Feito Y. Normative Values for Self-Reported Benchmark Workout Scores in CrossFit® Practitioners. *Sports Med Open.* 2018;4(1):39.
82. Prado Dantas TS, Aidar FJ, de Souza RF, de Matos GD, Pires Ferreira AR, de Almeida BN, Santos MDM, Barros GO, Santos CRR, da Silva Júnior WM. Evaluation of a CrossFit® Session on Post-Exercise Blood Pressure. *J Exerc Physiol Online.* 2018;21(1):44–51.
83. Ouellette KA, Brusseau TA, Davidson LE, Ford CN, Hatfield DL, Shaw JM, Eisenman PA. Comparison of the Effects of Seated, Supine, and Walking Interset Rest Strategies on Work Rate. *J Strength Cond Res.* 2016;30(12):3396–404.
84. de Sousa AF, dos Santos GB, dos Reis T, Valerino AJ, Del Rosso S, Boulosa DA. Differences in Physical Fitness between Recreational CrossFit® and Resistance Trained Individuals. *J Exerc Physiol Online.* 2016;19(5):112–22.
85. Fernández JF, Solana RS, Moya D, Marin JMS, Ramón MM. Acute physiological responses during crossfit® workouts. *European Journal of Human Movement.* 2015;35:114–24.
86. Butcher SJ, Neyedly TJ, Horvey KJ, Benko CR. Do physiological measures predict selected CrossFit® benchmark performance? *Open Access J Sports Med.* 2015;6:241–7.
87. Lundberg TR, Fernandez-Gonzalo R, Gustafsson T, Tesch PA. Aerobic exercise does not compromise muscle hypertrophy response to short-term resistance training. *J Appl Physiol (1985).* 2013;114(1):81–9.
88. Hickson RC. Interference of strength development by simultaneously training for strength and endurance. *Eur J Appl Physiol Occup Physiol.* 1980;45(2–3):255–63.
89. Berryman N, Mujika I, Bosquet L. Concurrent Training for Sports Performance: The 2 Sides of the Medal. *Int J Sports Physiol Perform.* 2019;14(3):279–85.
90. Wilson JM, Marin PJ, Rhea MR, Wilson SM, Loenneke JP, Anderson JC. Concurrent training: a meta-analysis examining interference of aerobic and resistance exercises. *J Strength Cond Res.* 2012;26(8):2293–307.
91. Kazior Z, Willis SJ, Moberg M, et al. Endurance Exercise Enhances the Effect of Strength Training on Muscle Fiber Size and Protein Expression of Akt and mTOR. *PLoS ONE.* 2016;11(2): e0149082.
92. Murach KA, Bagley JR. Skeletal Muscle Hypertrophy with Concurrent Exercise Training: Contrary Evidence for an Interference Effect. *Sports Med.* 2016;46(8):1029–39.
93. Buckner SL, Mouser JG, Jessee MB, Dankel SJ, Mattocks KT, Loenneke JP. What does individual strength say about resistance training status? *Muscle Nerve.* 2017;55(4):455–7.
94. Summitt RJ, Cotton RA, Kays AC, Slaven EJ. Shoulder Injuries in Individuals Who Participate in CrossFit Training. *Sports Health.* 2016;8(6):541–6.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.